

Performance, nutrient digestibility, and quantitative carcass traits of lambs fed sunflower seeds and vitamin E

Desempenho, digestibilidade dos nutrientes e características quantitativas da carcaça de cordeiros alimentados com grãos de girassol e vitamina E

Fabiana Alves de Almeida^{1*}; Américo Garcia da Silva Sobrinho²;
Gabriela Milani Manzi³; Natália Ludmila Lins Lima⁴; Ronaldo Oliveira Sales⁵;
Nivea Maria Brancacci Lopes Zeola²; Viviane Endo⁶; Thiago Henrique Borghi⁶

Abstract

The aim of this study was to evaluate performance, nutrient digestibility, and carcass quantitative traits of lambs fed diets with and without sunflower seeds and vitamin E. Thirty-two uncastrated lambs at approximately 60 days of age, with 15 ± 0.2 kg, were housed in individual stalls and slaughtered at 32.0 ± 0.2 kg body weight. Treatments consisted of the following four diets: C = sugarcane + concentrate; CS = sugarcane + concentrate with sunflower seeds; CE = sugarcane + concentrate with 1,000 mg vitamin E kg^{-1} diet dry matter (DM); and CSE = sugarcane + concentrate with sunflower seeds and 1,000 mg vitamin E kg^{-1} diet DM. The experimental design was completely randomized, in a 2×2 factorial arrangement (with and without sunflower seeds and with or without vitamin E, on the diet DM basis), and means were compared by Tukey's test at 5% significance level. Consumption of vitamin E intake by the lambs provided ($P < 0.05$) the highest ether extract intake (20.91 g day^{-1}) and consequently the highest ($P < 0.05$) digestibility of this nutrient (85.34%); however, it reduced ($P < 0.05$) the intakes of total carbohydrates ($394.81 \text{ g day}^{-1}$) and non-fibrous carbohydrates ($242.47 \text{ g day}^{-1}$) as a direct consequence of the diet composition. Inclusion of sunflower seeds and vitamin E in the lamb diets did not influence ($P > 0.05$) carcass morphological or quantitative measurements. Sunflower seeds and vitamin E showed to be satisfactory alternatives for inclusion in the concentrate for feedlot lambs, providing good performance and quality carcasses.

Key words: Antioxidant. Commercial cuts. Oilseeds. Sheep.

Resumo

O objetivo deste trabalho foi avaliar o desempenho, a digestibilidade dos nutrientes e as características quantitativas da carcaça de cordeiros submetidos a dietas contendo ou não grãos de girassol e vitamina

¹ Pós-Doutoranda, Departamento de Parasitologia, Instituto de Biociências, Universidade Estadual Paulista "Júlio de Mesquita Filho", UNESP, Botucatu, SP, Brasil. E-mail: faalvesalmeida@yahoo.com.br

² Profs., Departamento de Zootecnia, Faculdade de Ciências Agrárias e Veterinárias, UNESP, Jaboticabal, SP, Brasil. E-mail: americo@fcav.unesp.br; nivea.brancacci@ig.com.br

³ Pesquisadora, Scott Consultoria, Bebedouro, SP, Brasil. E-mail: gabmanzi@bol.com.br

⁴ Discente de Doutorado, Programa de Pós-Graduação em Zootecnia, Universidade Federal de Minas Gerais, UFMG, Belo Horizonte, MG, Brasil. E-mail: natludmila@yahoo.com.br

⁵ Prof., Departamento de Zootecnia, Universidade Federal do Ceará, UFC, Fortaleza, CE, Brasil. E-mail: ronaldosales@secrel.com.br

⁶ Discentes de Doutorado, Programa de Pós-Graduação em Zootecnia, Faculdade de Ciências Agrárias e Veterinárias, UNESP, Jaboticabal, SP, Brasil. E-mail: endo_vica@hotmail.com; thiagoborghi@zootecnista.com.br

* Author for correspondence

E. Foram utilizados 32 cordeiros machos não castrados com aproximadamente 60 dias de idade e $15 \pm 0,2$ kg, alojados em baias individuais e abatidos aos $32,0 \pm 0,2$ kg de peso corporal. Os tratamentos foram compostos por quatro rações: C- cana-de-açúcar + concentrado; CG- cana-de-açúcar + concentrado com grãos de girassol; CE- cana-de-açúcar + concentrado com 1000 mg vitamina E kg^{-1} de matéria seca (MS) da ração e CGE- cana-de-açúcar + concentrado com grãos de girassol e 1000 mg vitamina E kg^{-1} de MS da ração. O delineamento experimental foi o inteiramente casualizado em esquema fatorial 2×2 (com ou sem grãos de girassol e com ou sem vitamina E na base da MS da ração) e as médias comparadas pelo teste de Tukey a 5% de significância. O consumo de vitamina E pelos cordeiros melhorou o ganho médio diário de peso (0,241 kg). A inclusão dos grãos de girassol proporcionou ($P < 0,05$) maior consumo de extrato etéreo (20,91 g dia^{-1}) e conseqüentemente maior ($P < 0,05$) digestibilidade desse nutriente (85,34%), no entanto reduziu ($P < 0,05$) os consumos de carboidratos totais (394,81 g dia^{-1}) e não fibrosos (242,47 g dia^{-1}) como consequência direta da composição da ração. A inclusão de grão de girassol e vitamina E na dieta dos cordeiros não influenciou ($P > 0,05$) nas medidas morfológicas e quantitativas da carcaça. Grãos de girassol e vitamina E mostraram-se fontes alternativas satisfatórias para inclusão no concentrado para cordeiros confinados, proporcionando bom desempenho e carcaças com boa qualidade.

Palavras-chave: Antioxidante. Cortes comerciais. Grãos de oleaginosas. Ovinos.

Introduction

In intensive sheep meat production systems, aspects such as nutrition and dietary management are among the major factors responsible for increased productivity. The supply of diets of better quality reduces the finishing time for animals to reach slaughter weight, allowing the production of early lambs, with adequate fat cover, and thus meeting the consumer market demands (OLIVEIRA et al., 2016; SANTELLO et al., 2006). Because this production system is rather costly due to the high costs of inputs, the use of alternative ingredients in the diet, without impairing animal performance, is of great value.

Sugarcane is a forage plant commonly found on rural properties, especially in São Paulo state, that has been noteworthy for its lower production costs compared with other roughages, e.g., hay and corn silage (GALAN; NUSSIO, 2000). However, diets containing sugarcane require corrections in their protein and mineral contents, and a way to nutritionally complement these diets is by including oilseeds in their formulation (MORENO et al., 2010).

Oilseeds are the most widely used lipid sources in animal nutrition, usually replacing rapidly

fermentable carbohydrates, providing high energy density and viability to rumen fermentation and fiber digestion (TEIXEIRA; BORGES, 2005). However, oilseeds, such as sunflower seeds, may change the composition of the meat fatty acids, the unsaturated fatty acid content, and consequently increase the requirement of antioxidants like vitamin E for the preservation of meat products (DEMIREL et al., 2006; OLIVEIRA et al., 2011).

With liposoluble characteristics, the vitamin E accumulated in cell membranes protects the polyunsaturated fatty acids and other vulnerable components of cells from oxidative damage (BERCHIELLI et al., 2011). Their ingestion in the form of foods or supplements is also related to prevention of diseases, stimulation of the immune system, and modulation of aging-associated degenerative processes (BRIGELIUS-FLOHÉ et al., 2002).

The aim of this study was thus to evaluate animal performance, nutrient digestibility, and carcass quantitative traits of feedlot-finished lambs fed diets containing sugarcane as the roughage, associated with inclusion or lack of sunflower and vitamin E in the concentrate.

Material and Methods

This study complied with the principles of ethics in animal experimentation adopted by the Brazilian College of Animal Experimentation (*Colégio Brasileiro de Experimentação Animal*, COBEA), and was approved by the Committee for Ethics in Animal Use (*Comissão de Ética no Uso de Animais*, CEUA) of the Faculty of Agricultural and Veterinary Sciences (*Faculdade de Ciências Agrárias e Veterinárias*, FCAV – Unesp) under case no. 007670-09.

The experiment was conducted at the Faculty of Agricultural and Veterinary Sciences (*Faculdade de Ciências Agrárias e Veterinárias*, FCAV – Unesp), on Jaboticabal Campus, SP, Brazil (21°15'22" S and 48°18'58" W, 595 m altitude). Thirty-two newly weaned uncastrated male Ile de France lambs at approximately 60 days of age, with 15.0 ± 0.2 kg body weight (BW), were housed in individual stalls (covered shed) measuring approximately 1.0 m², with suspended slatted floor and equipped with feeder and drinker. At the onset of the experiment, lambs were identified, dewormed, vaccinated against clostridiosis, and distributed at random into the following treatments: C = sugarcane + concentrate; CS = sugarcane + concentrate with sunflower seeds; CE = sugarcane + concentrate with 1,000 mg vitamin E kg⁻¹ diet dry matter (DM); and CSE = sugarcane + concentrate with sunflower seeds and 1,000 mg vitamin E kg⁻¹ diet DM. The amount of 1,000 mg vitamin E (α -tocopherol acetate) was used per kg of diet DM, because, according to a study conducted by Lopez-Bote et al. (2001), this was the amount of vitamin that provided the best result.

The sugarcane, of variety IAC 86-2480, belonged to the experimental canebrake of FCAV and was harvested manually with a machete (on alternate days) and chopped (to 1 cm particles) in a stationary shredding machine, immediately before being supplied to animals. The concentrate was composed of ground corn, soybean meal, urea, sodium chloride, calcitic limestone, dicalcium

phosphate, and a mineral supplement. The roughage:concentrate ratio was 50:50, composing diets with similar protein (23.0%) and energy (3.0 Mcal metabolizable energy kg⁻¹ DM) contents. Experimental diets were formulated to meet the requirements of weaned lambs recommended by NRC (2007) for an average weight gain of 250 g day⁻¹ (Table 1).

Diets were supplied daily at 08h00 and 17h00, so as to allow 10% as leftovers (fresh matter); the weights of feed supplied and leftovers were recorded to estimate the dry matter intake (DMI). Lambs were weighed weekly to monitor and determine their average daily weight gain (ADG) and feed conversion (FC), obtained as the ratio between DMI (kg day⁻¹) and the average weight gain (kg). The period during which the animals were fed the diets varied with the feedlot period, which was 75 days, on average.

Samples of diets supplied and 10% of the leftovers from each animal were harvested weekly and stored in a freezer at -18 °C, with composite samples formed at the end of the experiment. These were pre-dried in a forced-air oven at 55 ± 5 °C, for 72 h, and later ground in a mill with knives and 1-mm sieve for determination of the DM, organic matter (OM), mineral matter (MM), crude protein (CP), and ether extract (EE), according to methods described by Silva and Queiroz (2006).

Concentrations of lignin, neutral detergent fiber corrected for ash and protein (NDFap), and acid detergent fiber (ADF) were determined according to Van Soest and Wine (1967). Total carbohydrates (TC) were calculated by the following equation: $TC = 100 - (\%CP + \%EE + \%MM)$, while non-fiber carbohydrates (NFC) were calculated as the difference between TC and NDFap, as proposed by Sniffen et al. (1992). The metabolizable energy content was calculated as 82% of the digestible energy.

Forty-five days after the beginning of the performance trial, the digestibility trial was started,

using 16 animals (four in each treatment) of the same 32 lambs, with an average body weight of 22.5 kg.

Lambs were housed in individual metabolic cages equipped with feeders and drinkers to determine feed intake and digestibility.

Table 1. Centesimal composition of ingredients and chemical composition of experimental diets.

Composition	Experimental diet ^a			
	C	CS	CE	CSE
Ingredient (% DM)				
Ground corn	7.93	0.22	7.90	0.22
Soybean meal	38.11	37.68	38.11	37.66
Sunflower seed	-	8.08	-	8.08
Sugarcane	49.92	50.10	49.92	50.00
Mineral mix ^b	0.49	0.49	0.49	0.49
Iodized salt	0.32	0.32	0.32	0.32
Dicalcium phosphate	0.81	0.81	0.81	0.81
Urea	1.33	1.14	1.33	1.15
Calcitic limestone	1.09	1.17	1.01	1.17
Vitamin E	-	-	0.10	0.10
Chemical composition ^c				
Dry matter ^d	43.08	43.17	43.08	42.98
Organic matter ^e	93.41	93.22	93.44	93.18
Mineral matter ^e	6.59	6.78	6.56	6.82
Crude protein ^e	23.59	23.36	23.59	23.37
Ether extract ^e	1.48	3.36	1.48	3.36
Lignin ^e	2.76	2.75	2.76	2.75
Neutral detergent fiber ^e	27.72	30.30	27.72	30.27
Acid detergent fiber ^e	14.31	16.71	14.31	16.69
Total carbohydrates ^{e,f}	68.34	66.23	68.37	66.45
Non-fibrous carbohydrates ^{e,g}	40.62	36.19	40.65	36.18
Metabolizable energy (Mcal kg ⁻¹ DM)	3.13	3.28	3.13	3.27

^aExperimental diet: C = sugarcane + concentrate; CS = sugarcane + concentrate with sunflower seeds; CE = sugarcane + concentrate with 1,000 mg vitamin E kg⁻¹ dietary dry matter; and CSE = sugarcane + concentrate with sunflower seeds and 1,000 mg vitamin E kg⁻¹ dietary DM.

^bMineral mix: phosphorus 50 g; calcium 150 g; sodium 100 g; magnesium 5 g; sulfur 25 g; zinc 1,500 mg; manganese 500 mg; cobalt 10 mg; iodine 40 mg; selenium 10 mg.

^cAnalyses performed at the Laboratory of Animal Nutrition (LANA) at FCAV – Unesp.

^d%.

^e% DM.

^fTotal carbohydrates = 100 – (%CP + %EE + %MM).

^gNon-fibrous carbohydrates = total carbohydrates – neutral detergent fiber.

^hMcal kg⁻¹ DM.

Lambs remained 12 days in the cages: seven for adapting and five for total feces collection, controlling the feed intake. The daily amount of feces excreted was collected and weighed, and 10% of the total were stored in a freezer at –18 °C. At the end of the collection period, samples were mixed to generate one composite sample per animal. The

intake of nutrients was calculated as the difference between the amount of a given nutrient present in the feed supplied and the amount of that nutrient in the leftovers. Subsequently, the intake of each digestible nutrient was calculated by multiplying the amount of a given nutrient consumed by its digestibility, expressed in g animal⁻¹ day⁻¹.

The digestibility coefficients of DM, OM, CP, NDFap, ADF, TC, and NFC were calculated as the difference between the amounts ingested and excreted in the feces, using the following formula: Apparent digestibility (%) = ((nutrient ingested, in g – nutrient excreted in the feces, in g)/nutrient ingested, in g) × 100.

Upon reaching 32.0 ± 0.2 kg body weight (approximately 135 days of age), lambs were weighed and fasted for solids for 16 h. Prior to slaughter, animals were weighed again to determine body weight at slaughter (BWS) and fasted weight (FW). Lambs were stunned by electronarcosis with a 250 V electric shock for two seconds, using an Imafrig Ltd. Futura stunning device, and then the jugular veins and carotid arteries were sectioned for bleeding, respecting the procedures that characterize humane slaughter. After skinning and evisceration, the gastrointestinal tract content (measured as the difference between full and empty gastrointestinal tract) was used in the determination of the empty body weight (EBW = BWS – gastrointestinal content – gallbladder content – bladder content) and dressing percentage (DP = (HCW – EBW)*100).

After evisceration, carcasses were weighed, generating the hot carcass weight (HCW), which was used to determine the hot carcass yield (HCY = HCW/BWS × 100), and transferred to a cold room at 4 °C, where they remained 24 h hung by the *Gastrocnemius* tendons on appropriate hooks, spaced 17 cm apart by the extremities. After this period, the cold carcass was weighed (CCW), so the chilling loss could be calculated (CL = (HCW – CCW/HCW) × 100). Conformation, which considers the distribution of muscle masses on the bone base, was evaluated subjectively always by the same person, according to Colomer-Rocher et al. (1988). Next, carcass morphological measurements were taken according to Osório et al. (1998), and the carcass compactness index ((CCI, kg/cm) = CCW/CIL (carcass internal length)) and leg compactness index (LCI = rump width/leg length) were calculated.

Later, the carcasses were divided lengthwise, and the left half was sectioned into five anatomical regions – neck, shoulder, ribs, loin, and leg – following Silva Sobrinho (2006), which were weighed individually to determine the representativeness in percentage values.

Measurements were taken in the dorsal portion of the *Longissimus lumborum* muscle, at the 12th and 13th thoracic ribs, to determine the loin eye area (LEA), calculated by the formula $(A/2 \times B/2)\pi$, proposed by Silva Sobrinho et al. (2003), where A is the maximum length and B is the maximum depth of the muscle, in cm. Additionally, measurements C (fat cover thickness) and GR (maximum cover thickness over the side part of the loin at 11 cm from the midline, in cm) were obtained using a digital caliper and a tape measure.

The experimental design was completely randomized, in a 2 × 2 factorial arrangement (with and without sunflower seeds and with or without vitamin E, on the diet DM basis). Data were subjected to analysis of variance by the F test at 5% significance level. When significant differences were detected between treatments for the different variables studied, they were compared by Tukey's test at the same significance level.

Results and Discussion

Lambs that remained on average 75 days confined consumed 793.43 g dry matter per day, which represented 3.36% of their body weight (BW) and a feed conversion of 3.44 kg DM intake kg⁻¹ weight gain. There was no interaction (P>0.05) between sunflower seeds and vitamin E for the analyzed variables, or differences (P>0.05) between treatments for initial body weight (IBW), final body weight (FBW), body weight at slaughter (BWS), days in the feedlot (DF), dry matter intake per day and per metabolic weight (DMI), and feed conversion (FC).

Dry matter intakes expressed in g animal⁻¹ day⁻¹, unit of metabolic size (g kg^{-0.75} day⁻¹), and percentage of body weight (% BW) were not influenced (P>0.05) by treatments, which may be because the diets had similar NDF contents and protein and energy values (Table 1). The same response was reported by Homem Junior et al. (2010), who did not observe differences in DM intake adding lipid sources to diets for Santa Ines lambs.

However, the average daily weight gain (ADG) was higher (P<0.05) in treatments with addition of

vitamin E (Table 2). The NRC (2007) recommends 15 to 30 mg vitamin E kg⁻¹ DM for growing sheep; however, it reports that levels above these, especially for growing animals, may improve the animal performance, as was observed in the present study. In a similar experiment, Wulf et al. (2003) added 1,000 mg vitamin E day⁻¹ to the diet of castrated lambs and observed a positive effect on weight gain. Macit et al. (2003) supplemented 45 mg day⁻¹ vitamin E in diets for Awassi sheep and reported a 6.7% improvement in daily weight gain.

Table 2. Performance of lambs fed diets with and without sunflower seeds and vitamin E.

Variable	Sunflower seeds (S)		Vitamin E (E)		S	E	Interaction (S × E)	CV (%)
	With	Without	With	Without	p	p	p	
IBW (kg)	15.04	14.99	15.03	15.00	0.570	0.780	0.610	1.32
FBW (kg)	32.24	32.09	32.26	32.07	0.330	0.190	0.910	1.01
BWS (kg)	30.28	30.19	30.20	30.28	0.670	0.690	0.860	1.57
DF (days)	76.90	72.80	71.90	77.80	0.210	0.080	0.550	9.47
DMI (g day ⁻¹)	770.55	816.32	800.42	786.44	0.080	0.580	0.800	7.06
DMI (g kg ^{-0.75} day ⁻¹)	71.87	76.38	74.64	73.60	0.060	0.650	0.790	6.93
DMI (%BW)	3.26	3.47	3.39	3.34	0.060	0.670	0.780	6.89
ADG (kg)	0.225	0.237	0.241 ^a	0.221 ^b	0.150	0.030	0.650	8.31
FC (DMI WG ⁻¹)	3.44	3.43	3.30	3.57	0.930	0.060	0.670	8.88

^{a,b}Means followed by different letters in the row differ by Tukey's test at 5% probability level; CV = coefficient of variation; IBW = initial body weight; FBW = final body weight; BWS = body weight at slaughter; DF = days in the feedlot; DMI = dry matter intake; BW = body weight; ADG = average daily weight gain; FC = feed conversion; WG = weight gain;

Crude protein intake did not differ (P>0.05) between treatments, averaging 157.68 g day⁻¹ (Table 3). Likewise, the intakes of OM, NDF, and ADF were not influenced (P>0.05) by addition of sunflower seeds and vitamin E to the diets. The similarity between treatments for intake was probably caused by the similar NDF contents and protein and energy values of the diets (Table 1).

Addition of sunflower seeds to lamb diets, irrespective of the use of vitamin E, influenced (P<0.05) the intakes of EE, TC, and NFC (Table 3). Ether extract intake (20.91 g day⁻¹) by the lambs from the treatment with sunflower seeds was higher than the others (9.54 g day⁻¹), which is a direct impact of the concentration of this nutrient in the diet (Table 1).

The intakes of TC and NFC of the lambs decreased from 459.46 and 304.42 g day⁻¹ when they received diets without inclusion of sunflower seeds to 394.81 and 242.47 g day⁻¹ with the diets containing the ingredient. Corn is the main quantitative source of carbohydrates in the diet of production animals, and when it is replaced with sunflower seeds, which contain lower amounts of these nutrients in their composition, the concentration of carbohydrates decreases, and consequently so does their intake. The intakes of EE, TC, and NFC observed in this study corroborate those obtained by Homem Junior et al. (2010), who evaluated inclusion of sunflower seeds in diets for feedlot Santa Ines lambs and reported higher EE intake and lower TC and NFC intakes when the seeds were added to the diets.

Table 3. Voluntary intake of nutrients (g) and gross energy (Mcal) obtained in the digestibility trial of lambs fed diets with and without sunflower seeds and vitamin E.

Variable (g/day)	Sunflower seeds (S)		Vitamin E (E)		S	E	Interaction (S × E)	
	With	Without	With	Without	p	p	p	CV (%)
DM	770.555	816.32	800.42	786.44	0.086	0.585	0.806	7.06
OM	504.322	565.49	543.75	526.06	0.090	0.609	0.835	14.17
CP	149.28	166.07	159.37	155.98	0.106	0.733	0.760	13.90
EE	20.91 ^a	9.54 ^b	15.09	15.37	0.001	0.798	0.919	15.89
NDF	152.78	155.04	157.30	150.51	0.851	0.574	0.816	17.19
ADF	65.92	56.68	61.65	60.96	0.149	0.911	0.627	22.22
TC	394.81 ^b	459.46 ^a	435.90	428.37	0.029	0.526	0.885	14.17
NFC	242.47 ^b	304.42 ^a	279.04	267.85	0.001	0.487	0.965	12.87

^{a,b}Means followed by different letters in the row differ by Tukey's test at 5% probability level; CV = coefficient of variation; DM = dry matter; OM = organic matter, CP = crude protein; EE = ether extract; NDF = neutral detergent fiber; ADF = acid detergent fiber; TC = total carbohydrates; NFC = non-fibrous carbohydrates.

The digestibility coefficients of DM and nutrients were similar ($P>0.05$) with and without inclusion of sunflower seeds and vitamin E, except for the variables EE and NFC, which were altered ($P<0.05$) with the inclusion of sunflower seeds (Table 4). There was an increase ($P<0.05$) from 68.46 to 85.34% in the digestibility coefficient of EE when the animals were fed diets containing sunflower seeds, which was probably due to the higher ether extract content of these diets (Table 1). Silva et al. (2007) stressed that lipid sources are good substitutes for the fermentable carbohydrates originating from corn because they have a high energy density, even though their inclusion in lamb diets reduces the digestibility of NFC. In the present study, the digestibility coefficient of NFC decreased ($P<0.05$) from 93.07 to 91.28%, probably due to the lower NFC intake (Table 3) by the animals fed diets with sunflower seed, affecting their use.

Table 5 shows the results for the carcass morphological measurements, which were not

influenced ($P>0.05$) by the treatments. The fact that there was no significance in the carcass measurements may be related to the standardization of slaughter according to body weight, since animals of the same breed and age, with similar body weights, show similar carcass morphological measurements. Similarly, the carcass compactness index ($0.24 \text{ kg}^{-1} \text{ cm}$) and leg compactness index (0.64), as well as the conformation (2.55) and fatness (2.40) scores were not affected ($P>0.05$) by treatments. These values corroborate Macedo et al. (2008), who evaluated the morphological measurements of the carcass of Suffolk crossbred lambs fed diets containing up to 19.80% sunflower seeds and slaughtered with 28 kg BW and did not observe differences in conformation (2.78), fatness (2.75), or compactness indices of carcass (0.27 kg cm^{-1}) and leg (0.63 kg cm^{-1}). Macit et al. (2003) and Homem Junior et al. (2010) also did not observe differences in morphological measurements of carcasses of lambs consuming vitamin E and sunflower seeds, respectively.

Table 4. Digestibility coefficients of nutrients and gross energy obtained in the digestibility trial of lambs fed diets with and without sunflower seeds and vitamin E.

Variable (%)	Sunflower seeds (S)		Vitamin E (E)		S	E	Interaction (S × E)	
	With	Without	With	Without	p	p	p	CV (%)
DC-DM	77.36	76.50	77.20	76.67	0.485	0.663	0.816	3.09
DC-OM	80.06	80.28	80.51	79.82	0.872	0.613	0.735	3.31
DC-CP	79.69	83.66	81.83	81.52	0.103	0.895	0.920	5.49
DC-EE	85.34 ^a	68.46 ^b	77.08	76.71	0.001	0.866	0.794	5.53
DC-NDF	35.13	31.37	33.37	33.13	0.297	0.947	0.7946	20.72
DC-ADF	27.80	26.91	27.11	27.60	0.694	0.829	0.915	16.11
DC-TC	73.11	73.98	73.69	73.40	0.648	0.876	0.577	5.09
DC-NFC	91.28 ^b	93.07 ^a	92.26	92.09	0.001	0.789	0.317	1.37
DC-GE	83.11	79.27	83.93	79.44	0.147	0.147	0.841	6.09

^{a,b}Means followed by different letters in the row differ by Tukey's test at 5% probability level; CV = coefficient of variation; DC = digestibility coefficient; DM = dry matter; OM = organic matter, CP = crude protein; EE = ether extract; NDF = neutral detergent fiber; ADF = acid detergent fiber; TC = total carbohydrates; NFC = non-fibrous carbohydrates; GE = gross energy.

Table 5. Morphological measurements of the carcass of lambs fed diets with and without sunflower seeds and vitamin E.

Variable	Sunflower seeds (S)		Vitamin E (E)		S	E	Interaction (S × E)	
	With	Without	With	Without	p	p	p	CV (%)
Conformation ^a	2.50	2.60	2.50	2.60	0.56	0.56	0.56	14.87
Fatness ^b	2.30	2.55	2.35	2.50	0.15	0.37	0.76	15.29
CEL (cm)	53.80	54.80	53.80	54.80	0.20	0.20	0.79	3.12
CIL (cm)	56.20	55.70	56.50	55.40	0.24	0.58	0.16	3.60
LL (cm)	32.80	34.30	33.30	33.80	0.27	0.71	0.94	8.89
CC (cm)	66.55	67.25	67.00	66.80	0.32	0.77	0.32	2.31
TC (cm)	37.20	36.15	36.50	36.85	0.43	0.79	0.39	8.04
RC (cm)	60.75	61.05	60.50	61.30	0.58	0.15	0.58	1.98
RW (cm)	21.77	21.80	21.69	21.88	0.97	0.82	0.68	8.70
CW (cm)	24.01	23.65	23.85	23.81	0.41	0.92	0.14	9.99
CD (cm)	24.14	24.23	23.96	24.41	0.79	0.21	0.55	3.19
CCI (kg cm ⁻¹)	0.24	0.24	0.23	0.25	0.84	0.75	0.13	6.68
LCI	0.66	0.63	0.65	0.64	0.64	0.52	0.87	13.54

Means followed by different letters in the row differ by Tukey's test at 5% probability level; ^aScore of 1 to 5, where 1= very poor and 5= excellent. ^bScore of 1 to 5, where 1 = absent fat and 5 = excess fat; CV = coefficient of variation; CEL = carcass external length; CIL = carcass internal length; LL = leg length; CC = chest circumference; TC = thigh circumference; RC = rump circumference; RW = rump width; CW = maximum chest width; CD = chest depth; CCI = carcass compactness index; LCI = leg compactness index.

Considering the mean values for carcass fatness, the higher ether extract contents stemming from inclusion of sunflower in the diets (Table 1) was assumed to result in greater fat deposition, which was not observed. The carcass compactness index

of 0.24 was similar to those found in the literature for lambs, with values ranging from 0.24 to 0.29 (ZUNDT et al., 2006).

According to Silva and Pires (2000), the quantitative evaluation of the carcass has a relevance

in the judgment of the performance attained by the animal during its development, in addition to being fundamental to the production process and relating to the end product, since it estimates the amount of meat produced available for sale. There was no difference ($P>0.05$) in the variables analyzed in the

animal carcass (Table 6). Results found for HCW and CCW were similar to those obtained by Macedo et al. (2008), who reported 13.89 kg HCW and 13.53 kg CCW in Suffolk lambs fed diets containing different levels of sunflower seeds and slaughtered with 28 kg BW.

Table 6. Quantitative carcass traits and percentages of cuts from the left half-carcass of lambs fed diets with and without sunflower seeds and vitamin E.

Variable	Sunflower seeds (S)		Vitamin E (E)		S	E	Interaction (S × E)	CV (%)
	With	Without	With	Without				
HCW (kg)	13.73	14.07	13.67	14.13	0.200	0.094	0.604	4.14
CCW (kg)	13.31	13.39	13.10	13.69	0.536	0.058	0.795	4.80
HCY (%)	45.32	46.60	45.27	46.66	0.171	0.139	0.672	4.34
CCY (%)	43.92	44.67	43.39	45.20	0.455	0.082	0.846	4.92
DP (%)	54.16	54.93	54.74	54.35	0.435	0.686	0.386	3.96
CL (%)	3.13	4.10	4.12	3.11	0.382	0.364	0.665	66.75
LHCW (kg)	6.61	6.84	6.50	6.95	0.560	0.063	0.834	4.75
Neck (%)	8.32	7.89	7.85	8.20	0.270	0.505	0.715	14.22
Shoulder (%)	20.88	20.47	20.77	20.58	0.463	0.765	0.998	7.20
Ribs (%)	25.11	25.58	25.38	25.32	0.581	0.825	0.296	5.99
Loin (%)	10.74	10.96	10.92	10.79	0.621	0.974	0.211	10.31
Leg (%)	35.25	34.65	35.23	34.68	0.352	0.345	0.159	4.53

^{a,b}Means followed by different letters in the row differ by Tukey's test at 5% probability level; CV = coefficient of variation; HCW = hot carcass weight; CCW = cold carcass weight; HCY = hot carcass yield; CCY = cold carcass yield; DP = dressing percentage; CL = chilling loss; LHCW = left half-carcass weight.

The cold carcass yield, which, according to Sañudo and Sierra (1986), can be influenced by the feeding system, ranged from 43.39 to 45.20% and remained close to the 44.5% recommended by Silva Sobrinho (2006). According to Grande et al. (2009), the chilling loss corresponds to the difference in weight before and after the carcass is chilled, and depends mainly on the conformation, fatness, and moisture loss. Thus, the larger the fat cover, the lower the chilling losses, given the increased protection provided to the carcass. The carcass chilling loss observed here (3.61%) was higher than the 1.94% obtained by Macedo et al. (2008). The mean value of 1.91 cm found for the fat cover of the carcasses (Table 7) was likely responsible for this greater weight loss from chilling.

Cuts' yields were 8.07% for neck, 20.70% for shoulder, 25.35% for ribs, 10.85% for loin, and 34.95% for leg (Table 6). The percentage evaluations of the different carcass cuts help in the selection of breeds and/or genetic groups that produce higher proportions of valued edible cuts and enable a quantitative assessment, as the carcass should have the best proportion possible of cuts with greater participation of muscles. The leg is the commercial cut of highest value in sheep carcass, and thus it is interesting that its proportion in the carcass be higher. Results for neck (8.06%), shoulder (20.68%), and leg (24.95%) yields obtained in the present study were similar to those observed by Macedo et al. (2008), who analyzed the carcass of Suffolk crossbred lambs fed diets containing up

to 19.80% sunflower seeds and reported yields of 7.67% for neck, 20.84% for shoulder, and 34.35%

for leg. According to Ortiz et al. (2005), in studies with lambs for meat production, these variables usually do not vary largely.

Table 7. Measurements of the *Longissimus lumborum* muscle of lambs fed diets with and without sunflower seeds and vitamin E.

Variable	Sunflower seeds (G)		Vitamin E (E)		S	E	Interaction (S × E)	
	With	Without	With	Without	P	P	P	CV(%) (%)
Measurement A (cm)	5.69	5.47	5.52	5.64	0.531	0.726	0.622	13.25
Measurement B (cm)	2.65	2.66	2.69	2.62	0.961	0.701	0.699	15.26
Measurement C (cm)	1.85	1.96	1.91	1.90	0.580	0.956	0.214	23.11
Measurement GR (cm)	2.06	2.25	2.06	2.26	0.498	0.427	0.474	25.67
LEA (cm ²)	11.94	11.59	11.71	11.82	0.531	0.257	0.050	15.79

^{a,b}Means followed by different letters in the row differ by Tukey's test at 5% probability level; CV = coefficient of variation; Measurement A = maximum length of the *Longissimus lumborum* muscle; Measurement B = maximum depth of the *Longissimus lumborum* muscle; Measurement C = minimum fat thickness over the *Longissimus lumborum*; Measurement GR = maximum fat thickness over the side part of the loin at 11 cm from the midline; LEA = loin-eye area, obtained by the formula $(A/2 \times B/2)\pi$.

Inclusion of sunflower seeds and vitamin E in the lamb diets did not influence ($P>0.05$) measurements A, B, C, or GR of the *Longissimus lumborum* muscle of the lambs (Table 7). The loin eye area obtained by the formula $(A/2 \times B/2)\pi$ proposed by Silva Sobrinho et al. (2003) was 11.76 cm², and is in line with the recommended range (8 to 14 cm²) for lambs slaughtered weighing 15 to 40 kg (SILVA SOBRINHO, 2006). This value was higher than the 10.30 cm² observed by Homem Junior et al. (2015), who evaluated the influence of supplementation with sunflower seeds on the LEA of the *Longissimus* muscle of lambs. The minimum fat thickness (measurement C) was below the minimum of 3 cm recommended by Silva Sobrinho (2006) for sheep carcasses; i.e., all treatments resulted in deficient fattening.

Measurements A and B of the *Longissimus* muscle have been studied by many researchers for being highly correlated with the proportion of muscle in the sheep carcass, besides LEA (PINHEIRO et al., 2009). Measurements A and B (5.58 and 2.65 cm) were similar to the 5.32 and 2.86 cm respectively, observed by Garcia et al. (2003), taken at the *L. dorsi*

muscle of Suffolk crossbred lambs slaughtered with 31 kg BW, and by Siqueira and Fernandes (2000), who evaluated the carcass of Corriedale lambs slaughtered with 31 kg and reported values of 5.10 and 2.40 for the same measurements.

The lipid level in the diets was low (Table 1), which likely influenced the lack of differences observed between treatments for the quantitative variables of carcass and commercial cuts. Nevertheless, despite the low lipid level of the diets, Sales et al. (2013) and Almeida et al. (2014), evaluating the qualitative and nutritional aspects, respectively, of meat from the same animal used in this experiment, reported benefits from a diet with inclusion of sunflower seeds and vitamin E on the meat water-holding capacity and shear force, as well as an improvement in fatty acid profile, with an increase in conjugated linoleic acid (CLA), among others.

Conclusions

Sunflower seeds and vitamin E are satisfactory alternative sources that can be included in the

concentrate for feedlot lambs, providing good performance and quality carcasses.

Acknowledgments

The authors thank the São Paulo Research Foundation (*Fundação de Amparo a Pesquisa do Estado de São Paulo*, FAPESP) for the financial support and the National Council for Scientific and Technological Development (*Conselho Nacional de Desenvolvimento Científico e Tecnológico*, CNPq) for the fellowship granted to the first author.

References

- ALMEIDA, F. A.; SILVA SOBRINHO, A. G.; MANZI, G. M.; LIMA, N. L. L.; ENDO, V.; ZEOLA, N. M. B. L. Dietary supplementation with sunflower seeds and vitamin E for fattening lambs improves the fatty acid profile and oxidative stability of the *Longissimus lumborum*. *Animal Production Science*, Melbourne, v. 55, n. 8, p. 1030-1036, 2014.
- BERCHIELLI, T. T.; PIRES, A. V.; OLIVEIRA, S. G. *Nutrição de ruminantes*. 2. ed. Jaboticabal: Funep, 2011. 616 p.
- BRIGELIUS-FLOHÉ, R.; KELLY, F. J.; SALONEM, J. T.; NEUZIL, J.; ZINGG, J. M.; AZZI, A. The European perspective on vitamin E: current knowledge and future research. *American Journal of Clinical Nutrition*, Bethesda, v. 76, n. 4, p. 703-716, 2002.
- COLOMER-ROCHER, F.; DELFA, R.; SIERRA, I. Método normalizado para el estudio de los caracteres cuantitativos y cualitativos de las canales, según los sistemas de producción. In: _____. *Método normalizado para el estudio de los caracteres cuantitativos y cualitativos de las canales caprinas y ovinas*. Madrid: INIA, 1988. p. 19-41.
- DEMIREL, G.; OZPINAR, H.; NAZLI, B.; KESER, O. Fatty acids of lamb meat from two breeds fed different forage: concentrate ratio. *Meat Science*, Barking, v. 72, n. 2, p. 229-235, 2006.
- GALAN, V. B.; NUSSIO, L. G. Novos custos para cana-de-açúcar. In: GALAN, V. B. (Ed.). *Boletim do leite*. Piracicaba: CEPEA/FEALQ, 2000. n. 74, p. 1-4.
- GARCIA, C. A.; MONTEIRO, A. L. G.; COSTA, C.; NERES, M. A.; ROSA, G. J. M. Medidas objetivas e composição tecidual da carcaça de cordeiros alimentados com diferentes níveis de energia em *creep feeding*. *Revista Brasileira de Zootecnia*, Viçosa, MG, v. 32, n. 6, p. 1380-1390, 2003.
- GRANDE, P. A.; ALCALDE, C. R.; LIMA, L. S.; AYER, I. M.; MACEDO, F. A. F.; MATSUSHITA, M. Características quantitativas da carcaça e qualitativas do músculo Longissimus dorsi de cabritos $\frac{3}{4}$ Boer + $\frac{1}{4}$ Saanen confinados recebendo rações contendo grãos de oleaginosas. *Revista Brasileira de Zootecnia*, Viçosa, MG, v. 38, n. 6, p. 1104-1113, 2009.
- HOMEM JUNIOR, A. C.; EZEQUIEL, J. M. B.; FÁVARO, V. R.; PEREZ, H. L.; ALMEIDA, M. T. C.; PASCHOALOTO, J. R.; D'ÁUREA, A. P.; CARVALHO, V. B.; NOCERA, B. F. Fontes de lipídios e classe sexual no confinamento de ovinos. *Semina: Ciências Agrárias*, Londrina, v. 36, n. 3, p. 2165-2174, 2015.
- HOMEM JÚNIOR, A. C.; EZEQUIEL, J. M. B.; GALATI, R. L.; GONÇALVES, J. S.; SANTOS, V. C.; SATO, R. A. Grãos de girassol ou gordura protegida em dietas com alto concentrado e ganho compensatório de cordeiros em confinamento. *Revista Brasileira de Zootecnia*, Viçosa, MG, v. 39, n. 3, p. 563-571, 2010.
- LOPEZ-BOTE, C. J.; DAZA, A.; SOARES, M.; BERGES, E. Dose-response effect of dietary vitamin E concentration on meat quality characteristics in light-weight lambs. *Journal of Animal Science*, Champaign, v. 73, n. 1, p. 451-457, 2001.
- MACEDO, V. P.; GARCIA, C. A.; SILVEIRA, A. C.; MONTEIRO, A. L. G.; MACEDO, F. A. F.; SPERS, R. C. Composições tecidual e química do lombo de cordeiros alimentados com rações contendo semente de girassol em comedouros privativos. *Revista Brasileira de Zootecnia*, Viçosa, MG, v. 37, n. 10, p. 1860-1868, 2008.
- MACIT, M.; AKSAKAL, V.; EMSEN, E.; ESENBUGA, N.; İRFAN AKSU, M. Effects of vitamin e supplementation on fattening performance, non-carcass components and retail cut percentages, and meat quality traits of Awassi lambs. *Meat Science*, Barking, v. 64, n. 1, p. 1-6, 2003.
- MORENO, G. M. B.; SILVA SOBRINHO, A. G.; LEÃO, A. G.; LOUREIRO, C. M. B.; PEREZ, H. L.; ROSSI, R. C. Desempenho, digestibilidade e balanço de nitrogênio em cordeiros alimentados com silagem de milho ou cana-de-açúcar e dois níveis de concentrado. *Revista Brasileira de Zootecnia*, Viçosa, MG, v. 39, n. 4, p. 853-860, 2010.
- NATIONAL RESEARCH COUNCIL – NRC. Nutrient requirements of sheep. Washington: National Academy Press, 2007. 408 p.
- OLIVEIRA, D. M.; LAEDIRA, M. M.; CHIZZOTTI, M. L.; MACHADO NETO, O. R.; RAMOS, E. M.; GONCALVES, T. M.; BASSI, M. S.; LANNA, D. P.

- D.; RIBEIRO, J. S. Fatty acid profile and quantitative characteristics of meat from Zebu steers fed with different oilseeds. *Journal of Animal Science*, Champaign, v. 89, n. 8, p. 2546-2555, 2011.
- OLIVEIRA, E. R.; MONÇÃO, F. P.; GABRIEL, A. M. A.; FERNANDES, A. R. M.; NASCIMENTO, F. A.; ABREU, F. S.; SILVA, L.V. M.; SILVA, L. H. P. Characteristics of carcasses and meat from feedlot lambs fed with sunflower cake. *Semina: Ciências Agrárias*, Londrina, v. 37, n. 1, p. 331-344, 2016.
- ORTIZ, J. S.; COSTA, C.; GARCIA, C. A.; SILVEIRA, L. A. Medidas objetivas das carcaças e composição química do lombo de cordeiros alimentados e terminados com três níveis de proteína bruta em *creep feeding*. *Revista Brasileira de Zootecnia*, Viçosa, MG, v. 34, n. 6, p. 2382-2389, 2005.
- OSÓRIO, J. C.; OSÓRIO, M. T.; JARDIM, P. O.; PIMENTEL, M. A.; POUHEY, J. L.; CARDELLINO, R. A.; MOTTA, L.; ESTEVES, R. *Métodos para avaliação da produção de carne ovina: in vivo, na carcaça e na carne*. Pelotas: Editora Universitária, UFPel, 1998. 107 p.
- PINHEIRO, R. S. B.; SILVA SOBRINHO, A. G.; SOUZA, H. B. A.; YAMAMOTO, S. M. Qualidade de carnes provenientes de cortes da carcaça de cordeiros e de ovinos adultos. *Revista Brasileira de Zootecnia*, Viçosa, MG, v. 38, n. 9, p. 1790-1796, 2009.
- SALES, R. O.; SILVA SOBRINHO, A. G.; ZEOLA, N. M. B. L.; LIMA, N. L. L.; MANZI, G. M.; ALMEIDA, F. A.; ENDO, V. Fresh and matured lamb meat quality fed with sunflower seeds and vitamin E. *Ciência Rural*, Santa Maria, v. 43, n. 1, p. 151-157, 2013.
- SANTELLO, G. A.; MACEDO, F. A. F.; MEXIA, A. A.; SAKAGUTI, E. S.; DIAS, F. J.; PEREIRA, M. F. Características de carcaça e análise do custo de sistemas de produção de cordeiras ½ Dorset ½ Santa Inês. *Revista Brasileira de Zootecnia*, Viçosa, MG, v. 35, n. 4, p. 1852-1859, 2006.
- SAÑUDO, C.; SIERRA, I. Calidad de la canal en la especie ovina. In: BUXADÉ, C. *Ovino*. Barcelona: Exclusivas ONE, 1986. p. 127-153.
- SILVA, D. J.; QUEIROZ, A. C. *Análise de alimentos: métodos químicos e biológicos*. 3. ed. Viçosa, MG: UFV, 2006. 235 p.
- SILVA, L. F.; PIRES, C. C. Avaliação quantitativa das proporções de osso, músculo e gordura da carcaça em ovinos. *Revista Brasileira de Zootecnia*, Viçosa, MG, v. 29, n. 4, p. 1253-1260, 2000.
- SILVA, M. M. C.; RODRIGUES, M. T.; RODRIGUES, C. A. F.; BRANCO, R. H.; RODRIGUES, C. A. F.; SARMENTO, J. L. R.; QUEIROZ, A. C.; SILVA, S. P. Suplementação de lipídios em dietas para cabras em lactação: consumo e eficiência de utilização de nutrientes. *Revista Brasileira de Zootecnia*, Viçosa, MG, v. 36, n. 1, p. 257-267, 2007.
- SILVA SOBRINHO, A. G. *Criação de ovinos*. 3. ed. Jaboticabal: Funep, 2006. 302 p.
- SILVA SOBRINHO, A. G.; KADIM, I. T.; PURCHAS, R. W. Effect of genotypes and age on carcass and meat quality characteristics of ram lambs. *Agricultural and Marine Science*, Corvallis, v. 8, n. 2, p. 73-78, 2003.
- SIQUEIRA, E. R.; FERNANDES, S. Efeito do genótipo sobre as medidas objetivas e subjetivas da carcaça de cordeiros terminados em confinamento. *Revista Brasileira de Zootecnia*, Viçosa, MG, v. 29, n. 1, p. 306-311, 2000.
- SNIFFEN, C. J.; CONNOR, J. D.; VAN SOEST, P. J. A net carbohydrate and protein system for evaluation cattle diets. II Carbohydrate and protein availability. *Journal of Animal Science*, Champaign, v. 70, n. 11, p. 3562-3577, 1992.
- TEIXEIRA, D. A. B.; BORGES, I. Efeito do nível de caroço de algodão sobre o consumo e a digestibilidade aparente da fração fibrosa do feno de braquiária (*Brachiaria decumbens*) em ovinos. *Arquivos Brasileiros de Medicina Veterinária e Zootecnia*, Belo Horizonte, v. 57, n. 2, p. 229-233, 2005.
- VAN SOEST, P. J.; WINE, R. H. Use of detergents in the analysis of fibrous feeds IV determination of plant cell wall constituents. *Journal Association Official Analytical Chemists*, Gaithersburg, v. 50, n. 1, p. 50-55, 1967.
- WULF, D. M.; MORGAN, J. B.; SANDERS, S. K.; TATUM, J. D.; SMITH, G. C.; WILLIAMS, S. Effects of dietary supplementation of vitamin e on storage and case life properties of lamb retain cuts. *Journal of Animal Science*, Champaign, v. 73, n. 2, p. 399-405, 2003.
- ZUNDT, M.; MACEDO, F. A. F.; ASTOLPHI, J. L. L.; MEXIA, A. A.; SAKAGUTI, E. S. Desempenho e características de carcaça de cordeiros Santa Inês confinados, filhos de ovelhas submetidas à suplementação alimentar durante a gestação. *Revista Brasileira de Zootecnia*, Viçosa, MG, v. 35, n. 3, p. 928-935, 2006.