

Growth and carcass characteristics of different crosses of broiler chickens reared under an alternative system

Características de crescimento e carcaça de diferentes cruzamentos de frangos criados em sistema alternativo

Fábio Loures Cruz^{1*}; Lorena Kathyryne Vieira Saraiva²; Grace Ellen Silva³; Tatiane Mendonça Nogueira⁴; Ana Paula Silva⁵; Peter Bitencourt Faria⁶

Abstract

This study aimed to assess the effect of different breeds and crosses of broiler chickens and sex on growth, carcass characteristics, and tissue composition. The experimental design was a completely randomized design in a 6×2 factorial scheme, with six genotypes (New Hampshire – NHS, Jersey Black Giant – JBG, Rhode Island Red – RIR, Indigenous Giant – IG, and the hybrids IG \times NHS and IG \times JBG) and two sexes (male and female). Each treatment was composed of five replications with three broiler chickens, totaling 180 broiler chickens. Zootechnical performance and growth were assessed through the Gompertz equation parameters. After slaughtering at 105 days, the characteristics of carcass and main cuts were determined by means of weight, yield, and proportion of tissue constituents (meat, bone, and skin). Regarding the Gompertz equation parameters, the genotype NHS presented a higher growth potential. Males presented higher live weight and carcass weights, except for IG. No difference was observed between genotypes and sexes ($P > 0.05$) for carcass yield. The genotype JBG presented the highest average for breast yield (24.4%). Males presented the highest average for leg yield (30.5%). The genotype IG presented the highest value of breast meat yield (71.2%), being similar to the genotypes from the crosses IG \times NHS, and IG \times JBG (68.9 and 68.3%, respectively). For meat to breastbone ratio, the genotype IG presented the highest value (3.4) and, considering the sex, females presented a higher average (3.2) for this characteristic. Regarding the proportion of leg meat, an effect was observed for genotype and sex ($P < 0.05$) with the highest values observed for the hybrid IG \times JBG and females (66.3 and 66.0%, respectively). For meat to leg bone ratio, the genotypes NHS, JBG, RIR, and IG \times NHS presented the highest values (2.6, 2.7, 2.6, and 2.6, respectively), with females presenting the highest value ($P < 0.05$) (2.7). The genotypes IG, IG \times NHS and IG \times JBG presented the best growth and carcass characteristics, being recommended to be reared under an alternative system. Females presented the highest breast yield and, proportionally, the highest amount of leg meat, which are important attributes in a chicken meat production system.

Key words: Breeds. Carcass yield. Gender. Indigenous Giant. Tissue constituents.

¹ Me. em Ciências Veterinárias, Discente de Doutorado, Universidade Federal de Lavras, UFLA, Lavras, MG, Brasil. E-mail: fabiolourescruz@gmail.com

² Médica Veterinária, Discente de Mestrado, Universidade Federal de Minas Gerais, UFMG, Montes Claros, Brasil. E-mail: lorena_kvs@hotmail.com

³ Discente, Curso de Graduação em Medicina Veterinária, UFLA, Lavras, MG, Brasil. E-mail: grace_ellens@hotmail.com

⁴ M.e, em Ciências Veterinárias, UFLA, Lavras, MG, Brasil. E-mail: tati.mn@hotmail.com

⁵ Médica Veterinária, E-mail: anysilvas27@hotmail.com

⁶ Prof. Adjunto, UFLA, Lavras, MG, Brasil. E-mail: peterbfvet@yahoo.com.br

* Author for correspondence

Resumo

Objetivou-se avaliar o efeito de diferentes raças e cruzamentos de frangos e do sexo sobre as características de crescimento, carcaça e composição dos tecidos. O delineamento foi inteiramente casualizado disposto em esquema fatorial (6x2), sendo seis genótipos (New Hampshire – NHS; Gigante Negra de Jersey – GNJ; Rodhe Island Red – RIR; Índio Gigante – IG; cruzamento entre as raças IG e NHS – IG x NHS e entre IG e GNJ – IG x GNJ) e dois sexos (macho e fêmea), com cinco repetições e três aves por repetição, totalizando 180 aves. Foi avaliado o desempenho zootécnico e o crescimento, através de parâmetros da equação de gompertz. Após o abate, que ocorreu aos 105 dias, foram determinadas as características de carcaça e dos principais cortes através do peso, do rendimento e da proporção dos constituintes teciduais (carne, osso e pele). Com relação aos parâmetros da equação de gompertz, as aves da raça NHS apresentaram maior potencial de crescimento. Os machos apresentaram médias superiores de peso vivo e de carcaça, exceto para o genótipo IG. Não houve diferença entre genótipos e sexos ($P > 0,05$) para o rendimento de carcaça. O genótipo GNJ apresentou a maior média de rendimento de peito (24,4 %). Para o rendimento de perna, os machos apresentaram a maior média (30,5 %). A raça IG apresentou maior valor de rendimento de carne do peito (71,2 %), sendo semelhante aos genótipos oriundos dos cruzamentos, IG x NHS e IG x GNJ (68,9 e 68,3 %, respectivamente). Para a relação carne/osso do peito, as aves da raça IG apresentaram o maior valor (3,4) e, com relação ao sexo, as fêmeas apresentaram média superior (3,2). Com relação à proporção de carne da perna, houve efeito do genótipo e do sexo ($P < 0,05$), sendo que o genótipo IG x GNJ e as fêmeas apresentaram maiores valores (66,3 e 66,0 %, respectivamente). Para a relação carne/osso da perna, os genótipos NHS, GNJ, RIR e IG x NHS apresentaram os maiores valores (2,6, 2,7, 2,6 e 2,6, respectivamente). As fêmeas apresentaram maior valor ($P < 0,05$) para esta variável (2,7). Os genótipos IG, IG x NHS e IG x GNJ apresentaram as melhores características de crescimento e de carcaça sendo preconizadas para a produção em sistema alternativo. As fêmeas apresentaram maior rendimento de peito e proporcionalmente maior quantidade de carne na perna, que são atributos importantes em um sistema de produção de carne de frango.

Palavras-chave: Constituintes teciduais. Índio Gigante. Raças. Rendimento de carcaça. Sexo.

Introduction

In recent years, there has been a significant growth of alternative systems for rearing broiler chickens due to the modern demand of consumers for natural products with more pronounced flavors. This tendency was accompanied by an evolution in the genetic area, with the emergence of new strains of free-range broiler chickens that present rusticity and good productive indices (DEL CASTILHO et al., 2013).

For profitability of this alternative poultry activity, the factors interfering with the most important carcass parameters for the market need to be taken into consideration, such as carcass, breast, and thigh yields, which are the most valued cuts and are directly influenced by genetics and sex (FARIA et al., 2010). Differences between sexes can occur due to the effect of sexual dimorphism related to the production of androgen hormones,

which are responsible for muscle anabolism (DEL CASTILHO et al., 2013). The choice of genetics is essential for the economic return of poultry activity and production planning since the growth rate of bird can influence yield and carcass composition (SARICA et al., 2014; VELOSO et al., 2014).

In this context, small and medium farmers have performed crosses between birds of different breeds and strains looking for characteristics such as rusticity, precocity, and mainly good yield indices of carcass and noble cuts for rearing broilers under extensive systems. However, the knowledge of the main characteristics of production, growth, and carcass is necessary for breed selection to be used in crosses of broiler chickens. Thus, the aim of this study was to assess the effect of different genotypes and sex on growth, performance, carcass characteristics, and tissue composition of broiler chickens recommended to be reared under an alternative system.

Material and Methods

The experiment was conducted in the Poultry Sector of the Department of Animal Science of the Federal University of Lavras (UFLA), Lavras, MG, Brazil, from July to October 2014. Breeds of broiler chickens used under alternative systems were selected for the experiment. The experimental design was a completely randomized design arranged in a factorial scheme (6×2) with six genotypes (New Hampshire – NHS, Jersey Black Giant – JBG, Rhode Island Red – RIR, Indigenous Giant – IG, F1 generation resulted from the cross between Indigenous Giant roosters and New Hampshire fowls – IG \times NHS, and F1 generation resulted from the cross between Indigenous Giant roosters

and Jersey Black Giant fowls – IG \times JBG) and two sexes (male and female), totaling 12 treatments. Each treatment consisted of five replications with three broiler chickens, totaling 15 broiler chickens per treatment and a total 180 broiler chickens (90 males and 90 females).

The diet for free-range broiler chickens was composed of three diet formulations considering their rearing phases: initial (1 to 30 days) without access to the grazing area, growth (31 to 55 days), and final (56 to 105 days) (Table 1). Thirty broiler chickens of each genotype were raised in an experimental unit of 90 m² with access to grazing, food, and water ad libitum.

Table 1. Composition and calculated values of experimental diets for free-range broiler chickens according to the rearing phase and age range in days.

Ingredient (kg)	Starter (1 to 30)	Growing (31 to 55)	(56 to 105)
Corn	57,91	63,69	68,54
Soybean Meal	31,48	25,94	24,03
Wheat Meal	6,81	7,01	4,23
Bicalcium Phosphate	1,59	1,36	1,31
Limestone	1,35	1,26	1,18
Common Salt	0,38	0,35	0,33
Mineral Premix ¹	0,10	0,10	0,10
Premix Vitamin ²	0,10	0,10	0,10
DL-Methionine 99%	0,20	0,14	0,13
L-Lisine 78%	0,03		
Choline Chloride 60%	0,05	0,05	0,05
Total (kg)	100	100	100
Calculated values			
Crude protein (%)	20,00	18,00	17,00
ME ³ (kcal/kg)	2800	2870	2940
Calcium (%)	1,00	0,90	0,85
Available phosphorus (%)	0,42	0,37	0,35
Sodium (%)	0,17	0,16	0,15
digM + C ⁴ (%)	0,74	0,64	0,61
Digestible Lysine (%)	0,96	0,81	0,76
Digestible tryptophan (%)	0,22	0,19	0,18
Crude fiber (%)	3,32	3,14	2,86

¹Mineral premix: Manganese, 75000 mg; zinc, 70000 mg; iron, 50000 mg; copper, 8500 mg; iodine, 1500 mg; cobalt, 200 mg.

²Vitamin premix: Vitamin A, 7000000 IU; vitamin D3, 2100000 IU; vitamin E, 50000 mg; vitamin K3, 2000 mg; vitamin B1, 2000 mg; vitamin B2, 4000 mg; vitamin B6, 3000 mg; vitamin B12, 3000 mg; niacin, 39800 mg; pantothenic acid, 15620 mg; folic acid, 1000 mg; selenium, 200 mg; biotin, 100 mg; antioxidant, 100000 mg. ³ME: Metabolizable energy. ⁴M+Cdig: methionine plus digestible cystine.

For the growth study, the parameters of the Gompertz equation were determined. The assessments of weight gain, feed intake, and feed conversion were performed according to the methodology of Del Castilho et al. (2013). Data acquisition was only descriptive since the rearing of broiler chickens was conducted in separate pickets according to the genotypes.

Broiler chickens were weighed, identified, fasted for 8 hours, and slaughtered by cervical dislocation followed by bleeding at 105 days of age. After evisceration, carcasses were individually packed in plastic bags and cooled at 5 °C for a period of 24 hours, following the distribution of treatments and repetitions. Subsequently, these carcasses were weighed for calculating the carcass and cut yields, as well as tissue composition, as Faria et al. (2011).

The data were analyzed by using the statistical program SISVAR®. The variables with responses of significant effects in the analysis of variance for treatments and/or interactions were submitted to the Tukey's test at 5% significance. All procedures described were approved by the Ethics Committee on Animal Use (CEUA) of the Federal University of Lavras (protocol number 017/14).

Results and Discussion

Broiler chickens from the F1 generation from the crosses between Indigenous Giant (IG) and Jersey

Black Giant (JBG) (IG × JBG) presented higher estimated values of live weight at maturity (W_m) (Table 2). The highest value of maturity rate (b) was observed for the genotype New Hampshire (NHS), indicating that this genotype presents a higher daily weight gain and hence a higher growth potential (Table 2). In addition, NHS presented the lowest age for the maximum growth rate (T^*), evidencing their higher precocity. As the age increases, growth rate slows and the moment at which the acceleration pattern changes characterizes the inflection point, corresponding to the T^* value (KESSLER, 2000). Parameter values of the Gompertz equation of all genotypes presented a high reliability, with R^2 values above 0.98 (Table 2).

A numerical difference was observed between genotypes for weight gain, feed intake, and feed conversion (Table 2). Higher values of weight gain were observed for broiler chickens from the genotypes NHS, IG × NHS, and IG × JBG. These results show that crossbreeding provided offspring with higher potentials of weight gain when compared to purebred birds used at crosses, which is mainly due to the hybrid vigor effect. In addition to being considered pure, the genotype NHS presents a great potential for lean tissue deposition in the carcass, which gives it a higher weight gain rate, being used in breeding programs that originated many current hybrids of broiler chickens (ALBINO et al., 2014).

Table 2. Parameters of performance and Gompertz equations for live weight of chicken broilers of different breeds and crosses.

Parameters	Genotypes					
	NHS	JBG	RIR	IG	IG x NHS	IG x JBG
P_m (g) ^{(1)(a)}	2637	2832	2343	2800	3211	3620
b (per day) ^{(2)(a)}	0,0222	0,0175	0,0201	0,0205	0,0212	0,0197
T^* (day) ^{(3)(a)}	61	86	63	70	73	79
R^2	0,9955	0,9878	0,9887	0,9993	0,9989	0,9998
Weight Gain (g/ave) ^(a1)	144,6	138,4	126,0	123,1	149,1	143,1
feed intake (g/ave) ^(a2)	652,4	649,3	611,5	632,0	637,7	629,9
feed conversion ^(a2)	4,1	3,8	3,6	4,1	4,0	3,5

⁽¹⁾Weight at maturity; ⁽²⁾maturity rate; ⁽³⁾age at which growth rate is maximal; Gompertz equation: $P_t = P_m \times \exp(-\exp(-b \times (\text{age} - t)))$; ^(a)Descriptive data; ⁽¹⁾Period: 1 to 105 days; ⁽²⁾Period: 36 to 105 days. Namely: NHS – New Hampshire; JBG – Jersey Black Giant; RIR – Rhode Island Red; IG – Indigenous Giant; IG × NHS – F1 generation resulted from the cross between Indigenous Giant and New Hampshire; IG × JBG – F1 generation resulted from the cross between Indigenous Giant and Jersey Black Giant.

Broiler chickens from the genotypes NHS and JBG presented the highest feed intake (Table 2) and, in general, a great potential of weight gain and deposition of lean tissue in the carcass (ALBINO et al., 2014). According to Santos et al. (2005), broiler chickens with genetic characteristics that provide higher growth rates, i.e. higher potentials of weight gain, present a higher feed intake due to their higher nutritional requirements to express their genetic potential.

The best values of feed conversion were observed for broiler chickens from the genotype Rodhe Island Red (RIR) and those from crosses between IG and JBG (IG × JBG) (Table 2). In general, accelerated-growth broiler chickens are more efficient at converting ingested food into body weight gain than slow-growing broiler chickens (GONZALES et al., 1998). However, broiler chickens with greater potential of weight gains such as those from the genotypes NHS and JBG presented worse feed conversion rates mainly due to higher feed intake values. Moreover, broiler chickens from crosses (IG × NHS and IG × JBG) presented better values of feed conversion when compared to purebred birds used at crosses such as IG, NHS, and JBG, which indicates a positive effect of crossing on the efficiency in converting the ingested food into weight gain.

An interaction between sex and genotype was observed for live weight, carcass, and legs (Table 3). In this case, males presented the highest averages for all genotypes, except for the genotype IG, which did not show difference between sexes (Table 4). Among males, the genotypes RIR and IG presented the lowest averages for these variables whereas this behavior was observed for the genotype RIR among females. In addition, the results were similar among the other genetic groups.

In general, this behavior is related to the sexual dimorphism (GONZALES; SARTORI, 2002; DEL CASTILHO et al., 2013), in which males present a higher growth potential due to their capacity of muscular deposition (FARIA et al., 2011; MITROVIC et al., 2011). Because the genotype

RIR has better characteristics for egg production, they have lighter carcasses whereas broiler chickens from the genotype IG present later sexual maturity, which may have led to the lowest values of these variables since their gonads were not yet developed (ABCIG, 2016). Higher values of live weight and carcass are characteristic of broiler chicken strains with higher growth potentials, which present a higher meat production capacity (SARICA et al., 2014; VELOSO et al., 2014).

No effect of genotype and sex were observed for carcass yield values (Table 3). This result is regardless of genetic potential or sex category; and it would be related to the proportional growth rate between broiler chickens of the different tissues and organs (FARIA et al., 2010; MITROVIC et al., 2011; VELOSO et al., 2014).

Male broiler chickens from the genotypes NHS, RIR, IG × NHS, and IG × JBG and female broiler chickens from the genotypes JBG and IG showed higher average values of breast weight. Among males, the genotypes RIR and IG presented the lowest average values of breast weight whereas, among females, the genotypes NHS and RIR presented the lowest values (Table 4). In general, broiler chickens from genotypes with a lower degree of breeding for meat production, such as RIR and IG, tend to show a less development of noble parts of the carcass, such as the breast.

The genotype JBG presented the highest average of breast yield when compared to NHS, however similar to the other genotypes (Table 4). These results indicated that the crosses provided individuals with similar characteristics of breast yield, which is associated with growth rate, muscle development, and proportion of red/white fibers (MADEIRA et al., 2006; FARIA et al., 2010; MIKULSKI et al., 2011; SARICA et al., 2014).

In general, females presented the highest average values of breast yield, while males presented the highest values of leg yields (Table 3). These results would be related to the effect of sexual dimorphism, as found in other studies (FARIA et al., 2011; DEL CASTILHO et al., 2013; SARICA et al., 2014).

Table 3. Parameters of carcass and cuts of broiler chickens reared under an alternative system.

Variable	Average Overall	Sex		P ⁽¹⁾			CV ⁽²⁾ (%)
		Male	Female	Genotype (G)	Sex (S)	SxG	
Live Weight (kg)	1,8	2,1a	1,6b	0,001*	0,001*	0,015*	8,1
Carcass Weight (Kg)	1,4	1,5a	1,2b	0,001*	0,001*	0,007*	10,9
CY ⁽³⁾ (%)	75,6	73,8a	75,2a	0,506 ^{ns}	0,318 ^{ns}	0,306 ^{ns}	7,2
Breast (g)	306,6	327,1b	302,9a	0,001*	0,001*	0,001*	10,7
Breast (%)	22,4	21,6b	23,8a	0,007*	0,001*	0,284 ^{ns}	8,0
Breast Meat (g)	207,1	220,9a	193,2b	0,001*	0,001*	0,018*	13,1
Breast Meat (%)	67,8	67,2a	68,4a	0,001*	0,117 ^{ns}	0,167 ^{ns}	4,2
Breast Skin (g)	29,0	29,7a	27,6a	0,001*	0,065 ^{ns}	0,830 ^{ns}	18,9
Breast Skin (%)	9,6	9,2b	10,0a	0,001*	0,015*	0,022*	12,7
Breast Bone (g)	68,5	76,7a	60,4b	0,004*	0,001*	0,684 ^{ns}	14,6
Breast Bone (%)	22,6	23,6a	21,6b	0,252 ^{ns}	0,008*	0,773 ^{ns}	12,2
Breast meat/bone	3,2	2,9b	3,2a	0,018*	0,013*	0,353 ^{ns}	15,1
Leg (g)	413,3	474,5a	352,1b	0,001*	0,001*	0,007*	9,5
Leg (%)	29,9	30,5a	29,3b	0,129 ^{ns}	0,001*	0,186 ^{ns}	4,6
Leg Meat (g)	262,6	300,0a	227,4b	0,001*	0,001*	0,005*	11,5
Leg Meat (%)	65,0	64,4b	66,0a	0,014*	0,001*	0,639 ^{ns}	2,6
Leg Skin (g)	35,3	38,7b	31,9a	0,001*	0,001*	0,217 ^{ns}	13,4
Leg Skin (%)	8,9	8,3b	9,4a	0,001*	0,001*	0,021*	7,2
Leg Bone (g)	106,1	126,3a	85,2b	0,001*	0,001*	0,131 ^{ns}	12,4
Leg Bone (%)	26,1	27,3a	24,6b	0,001*	0,001*	0,700 ^{ns}	6,1
Leg meat/bone	2,6	2,4b	2,7a	0,002*	0,001*	0,858 ^{ns}	8,9
Wing (g)	166,8	186,8a	146,8b	0,001*	0,001*	0,005*	9,2
Wing (%)	12,3	12,41a	12,25a	0,175 ^{ns}	0,539 ^{ns}	0,229 ^{ns}	8,1
Back (g)	238,4	268,6a	208,1b	0,001*	0,001*	0,001*	10,8
Back (%)	17,5	17,7a	17,3a	0,279 ^{ns}	0,326 ^{ns}	0,296 ^{ns}	9,6
EV ⁽⁴⁾ (g)	87,95	97,6a	78,3b	0,336 ^{ns}	0,001*	0,043*	7,4
EV ⁽⁴⁾ (%)	6,6	6,5a	6,6 ^a	0,001*	0,724 ^{ns}	0,060 ^{ns}	9,6
AF ⁽⁵⁾ (g)	15,2	13,8a	16,5a	0,001*	0,053 ^{ns}	0,192 ^{ns}	34,6
AF ⁽⁵⁾ (%)	0,85	0,7b	1,0a	0,001*	0,001*	0,001*	33,3

⁽¹⁾Tukey's test at 5% probability; Means followed by lowercase letters in the row indicate difference between sexes; ^{ns} Non-significant; *Significant at 5% probability; ⁽²⁾Coefficient of variation; ⁽³⁾Carcass yield; ⁽⁴⁾Edible viscera (liver + heart + gizzard); ⁽⁵⁾ Abdominal fat.

Table 4. Live weight and characteristics of carcass and main cuts of broiler chickens of different genotypes raised under an alternative system.

Variable	Sex	Genotype					
		NHS	JGB	RIR	IG	IGxNHS	IGxJBG
Live Weight (Kg)	Male	2,2aA	2,1aAB	1,8aC	1,9aBC	2,2aA	2,2aA
	Female	1,5bAB	1,6bAB	1,4bB	1,7aA	1,6bAB	1,7bA
	Average	1,9AB	1,9AB	1,6C	1,8B	1,9AB	2,0A
Carcass Weight (Kg)	Male	1,6aA	1,5aAB	1,3aB	1,3aB	1,7aA	1,7aA
	Female	1,1bAB	1,2bAB	0,9bB	1,3aA	1,2bAB	1,3bA
	Average	1,4AB	1,3ABC	1,2C	1,3ABC	1,5A	1,5A
CY ⁽¹⁾ (%)	Average	74,5A	72,2A	73,2A	74,7A	76,1A	76,3A
	Male	326,2aAB	351,8bA	274,7aB	270,8bB	373,0aA	365,9aA
Breast (g)	Female	251,8bBC	392,1aAB	228,4bC	322,9aA	303,7bAB	318,4bA
	Average	288,9CD	371,9ABC	251,6D	296,8BC	338,3AB	342,2A
Breast (%)	Average	21,2B	24,4A	22,0AB	22,4AB	23,3AB	22,8AB
	Male	498,3aA	482,2aA	381,8aB	405,4aB	529,4aA	549,6aA
Leg (g)	Female	328,8bAB	350,1bAB	292,8bB	381,3aA	366,0bAB	393,8bA
	Average	413,6BC	416,1BC	337,3D	393,4C	447,7AB	471,7A
Leg (%)	Average	29,9A	29,8A	29,3A	29,8A	29,5A	31,0A

Values with different superscripts in the same row or column are statistically different according to the Tukey's test at 5% probability; means followed by lowercase letters in the column indicate the difference between sexes; means followed by uppercase letters in the row indicate the difference between genotypes. Namely: NHS – New Hampshire; JGB – Jersey Black Giant; RIR – Rhode Island Red; IG – Indigenous Giant; IG × NHS – F1 generation resulted from the cross between Indigenous Giant and New Hampshire; IG × JBG – F1 generation resulted from the cross between Indigenous Giant and Jersey Black Giant. ⁽¹⁾Carcass yield.

An interaction between genotype and sex was observed for breast and leg meat weight (Table 3). In all genotypes, males presented the highest values for these characteristics, except for IG, in which no differences between sexes were found (Tables 5 and 6). Among males, crossbred broiler chickens (IG × NHS and IG × JBG) presented the highest averages of breast meat weight when compared to RIR birds whereas similar values were found for the other genotypes. Among females, the genotypes IG and those from crosses (IG × NHS and IG × JBG) presented higher averages in relation to the genotypes NHS and RIR (Table 5).

Considering the leg meat weight, the crosses IG × NHS and IG × JBG presented the highest averages among males, differing from the genotype of origin IG (Table 6). This shows a muscular increment with crossbreeding. However, this effect was not observed among females since no difference was observed in leg meat weight among origin genotypes (NHS, JBG, and IG) and those from crosses (IG × NHS and IG × JBG).

In general, crossbred broiler chickens (IG × NHS and IG × JBG), regardless of sex, presented higher values of breast and leg meat weight (Tables 5 and 6). This behavior indicates a greater deposition of muscle tissue due to protein anabolism and hybrid vigor associated with the sexual category and crosses between breeds (GONZALES; SARTORI, 2002).

Table 5. Weight and proportion of tissue constituents (meat, bone, and skin) of the breast of broiler chickens from different genotypes reared under an alternative system.

Variable	Sex	Genotype					
		NHS	JBG	RIR	IG	IGxNHS	IGxJBG
Meat (g)	Male	205,4aAB	226,8aAB	188,9aB	203,3aAB	250,2aA	250,9aA
	Female	154,5bB	192,3bAB	150,8bB	235,4aA	210,4bA	216,0bA
	Average	179,9BC	209,6AB	169,8C	219,4A	230,3A	233,4A
Meat (%)	Average	65,0C	66,9BC	66,6BC	71,2A	68,9AB	68,3AB
Bone (g)	Average	65,8AB	71,2AB	60,2B	64,0B	72,2AB	77,7A
Bone (%)	Average	23,9A	22,8A	23,4A	21,3A	21,6A	22,6A
Skin (g)	Average	30,1AB	32,1A	25,4AB	23,2B	31,9A	29,2A
Skin (%)	Male	9,6bA	9,8aA	9,4aA	8,0aA	9,3aA	9,1aA
	Female	12,7aA	10,8aAB	10,5aAB	7,1aC	9,7aB	9,1aBC
	Average	11,2A	10,3AB	9,9AB	7,6C	9,5B	9,1BC
Maet/bone	Average	2,7B	3,1AB	2,9AB	3,4A	3,2AB	3,1AB

Values with different superscripts in the same row or column are statistically different according to the Tukey's test at 5% probability; means followed by lowercase letters (ab) in the column indicate the difference between sexes; means followed by uppercase letters (AB) in the row indicate the difference between genotypes. Namely: NHS – New Hampshire; JBG – Jersey Black Giant; RIR – Rodhe Island Red; IG – Indigenous Giant; IG × NHS – F1 generation resulted from the cross between Indigenous Giant and New Hampshire; IG × JBG – F1 generation resulted from the cross between Indigenous Giant and Jersey Black Giant.

A significant genotype effect was observed in the proportion of breast meat (Table 3). In this sense, the genotype IG presented the highest average value for this characteristic, which was similar to the values found for crosses (IG × NHS and IG × JBG) (Table 5). In fact, IG broiler chickens present a large size and hence a higher proportion of breast meat. Due to the heterosis effect, broiler chickens from the F1 generation (IG × NHS and IG × JBG) also presented similar proportions. On the other hand, the genotype IG presented the lowest proportion of leg meat, but its crossing with the genotype JBG (IG × JBG) provided a significant improvement of this characteristic, with an average higher than that found for the genotype IG, but similar to the others genotypes (Table 6).

Broiler chickens from crosses between the genotypes IG and JBG (IG × JBG) presented a higher value of breastbone weight when compared to the genotypes RIR and IG, but similar to the other genotypes (Table 5). Similar results were obtained for legs in which the broiler chickens of this genotype presented the highest average of bone weight, however similar to IG and broiler chickens from a cross between IG and NHS (IG × NHS) (Table 6). Regarding the proportion of breastbone, no difference was observed between genotypes (Table 5). However, the genotype IG presented a higher value for the proportion of leg bone when compared to the other genotypes (Table 6), which confirms its breed pattern, i.e. a leg with long length and a well-developed bone structure (ABCIG, 2016).

Table 6. Weight and proportion of tissue constituents (meat, bone, and skin) of the leg of broiler chickens from different genotypes reared under an alternative system.

Variable	Sex	Genotype					
		NHS	JBG	RIR	IG	IGxNHS	IGxJBG
Meat (g)	Male	293,0aABC	310,5aAB	260,6aBC	247,4aC	340,1aA	348,8aA
	Female	204,5bA	224,6bA	188,2bB	249,4aA	237,2bA	260,4bA
	Average	248,8BC	267,6AB	224,4C	248,4BC	288,6AB	304,6A
Meat (%)	Average	64,6AB	65,5AB	65,2AB	63,6B	65,8AB	66,3A
Bone (g)	Average	97,6BC	101,3BC	88,6C	112,0AB	113,4AB	121,5A
Bone (%)	Average	25,1B	24,7B	25,4B	28,5A	25,7B	26,3B
Skin (g)	Average	39,0A	39,5A	31,8BC	30,5C	37,0AB	34,0ABC
Skin (%)	Male	9,2bA	9,2bA	8,7bAB	7,7aBC	8,1aABC	7,2aC
	Female	11,4aA	10,4aAB	10,0aBC	8,0aD	8,8aCD	7,7aD
	Average	10,3A	9,8AB	9,3B	7,8CD	8,5C	7,4D
Maet/bone	Average	2,6A	2,7A	2,6A	2,2B	2,6A	2,5AB

Values with different superscripts in the same row or column are statistically different according to the Tukey's test at 5% probability; means followed by lowercase letters (ab) in the column indicate the difference between sexes; means followed by uppercase letters (AB) in the row indicate the difference between genotypes. Namely: NHS – New Hampshire; JBG – Jersey Black Giant; RIR – Rodhe Island Red; IG – Indigenous Giant; IG × NHS – F1 generation resulted from the cross between Indigenous Giant and New Hampshire; IG × JBG – F1 generation resulted from the cross between Indigenous Giant and Jersey Black Giant.

The variation in the amount of certain tissue will influence the proportion of other tissues and, in situations of similar weight, birds with a higher amount of muscle tissue will indicate individuals with a higher potential for growth and meat production (SOGUNLE et al., 2013). This was also observed for females that presented a higher proportion of meat and a lower proportion of bone in the legs (Table 3). Males presented a higher weight and proportion of bone for both cuts (Table 3), which confirms their higher live weight and, consequently, a well-developed bone structure (MITROVIC et al., 2011; DEL CASTILHO et al., 2013).

A difference between genotypes was observed for breast and leg skin weight (Table 3). In general, the genotype IG presented lower values for these characteristics, not differing from the genotypes RIR and NHS (breast skin weight) and RIR and IG × JBG (leg skin weight) (Tables 5 and 6). Regarding the proportion of skin, an interaction was observed between genotype and sex for both cuts (Table 3). For breast, a difference was observed between sexes only for the genotype NHS, with higher values for females (Table 5). Among males, no difference was

observed between genotypes for the proportion of skin. However, among females, the lowest value was found for the genotype IG, which was similar to the hybrid IG × JBG.

Regarding the leg, a difference was observed for the proportion of skin between sexes only for the genotypes NHS, JBG, and RIR, with the females presenting the highest averages (Table 6). In general, the genotypes NHS and JBG presented higher values of the proportion of leg skin for both sexes. The differences in weight and proportion of skin would be related to the influence of steroid hormones, which exert effects in its different layers such as the hypoderm, causing an increase in thickness and weight due to an increased volume of adipocytes that invade the superficial fascia (ISOLA et al., 2013).

For meat to bone ratio in the breast, the genotype IG presented higher values when compared to those from the genotype NHS, but similar to the other genetic groups (Table 5). The lowest value for meat to bone ratio in the leg was observed for IG broiler chickens, but similar to IG × JBG (Table 6). This behavior indicates that the tissue deposition rates

were different between chickens and may occur independently between cuts. In addition, a better balance in relation to this proportion and tissue composition may be obtained by crosses between different breeds, aiming at achieving a maximum efficiency in muscle development.

Regarding sex, females presented better average values of meat to bone ratio for both cuts (Table 3). According to Gonzales and Sartori (2002), this behavior can be explained mainly by the characteristics of sexual dimorphism, in which females present a less-developed bone structure, which is evidenced by the low values of bone weight found in our study for females.

An interaction was observed between genotype and sex for wing and back weight (Table 3), in which males from all genotypes presented higher

values, except for IG, which did not present differences between sexes (Table 7). Among males, the genotypes RIR and IG presented the lowest averages for these variables. Among females, on the other hand, similar results were found, with the genotype RIR presenting the lowest values of wing and back weight (Table 7). These differences between genotypes are due to differences in the tissue growth rates, i.e. even when broiler chickens are slaughtered at the same age, slower-growing broiler chickens present lower bone development (FARIA et al., 2010, 2011; SARICA et al., 2014). Thus, because wing and back are composed mostly of bone tissue, lesser-developing broiler chickens, such as those from the genotypes RIR and IG, tend to have these cuts less heavy. However, neither genotype nor sex effects were observed for wing nor back yields (Table 7).

Table 7. Characteristics of secondary cuts, weight, and yield of edible viscera and abdominal fat of broiler chickens from different genotypes reared under an alternative system.

Variable	Sex	Genotype					
		NHS	JBG	RIR	IG	IGxNHS	IGxJBG
Wing (g)	Male	188,9aAB	193,8aAB	163,8aC	167,0aBC	204,4aA	203,4aA
	Female	142,7bAB	144,5bAB	127,7bC	163,6aA	147,6bA	154,7bA
	Average	165,8AB	169,2A	145,7B	165,3AB	175,9A	179,1A
Wing (%)	Average	12,1A	12,8A	12,7A	12,6A	12,0A	11,8A
Back (g)	Male	282,8aAB	278,2aAB	234,8aBC	213,4aC	297,9aA	304,4aA
	Female	211,1bAB	206,9bAB	167,3bB	225,1aA	217,5bA	220,4bA
	Average	247,0AB	242,5AB	201,1C	219,3BC	257,8A	262,4A
Back (%)	Average	18,0A	18,4A	17,4A	16,6A	17,6A	17,2A
EV ⁽¹⁾ (g)	Male	103,1aA	97,7aA	96,5aA	91,1aA	94,9aA	102,1aA
	Female	72,8bA	82,2bA	80,1bA	79,4bA	75,9bA	79,2bA
	Average	87,9A	89,9A	88,3A	85,3A	85,4A	90,6A
EV ⁽¹⁾ (%)	Average	6,4BC	6,8B	7,7A	6,5BC	5,9C	5,9C
AF ⁽²⁾ (g)	Average	31,4A	29,4A	10,2BC	2,3D	4,7CD	13,0B
	Male	1,2bA	1,3bA	0,6aB	0,1aC	0,2aBC	0,6aB
	Female	2,4aA	1,9aA	0,7aBC	0,2aC	0,4aBC	0,7aB
AF ⁽²⁾ (%)	Average	1,8A	1,6A	0,7B	0,1C	0,3C	0,7B

Values with different superscripts in the same row or column are statistically different according to the Tukey's test at 5% probability; means followed by lowercase letters (ab) in the column indicate the difference between sexes; means followed by uppercase letters (AB) in the row indicate the difference between genotypes. Namely: NHS – New Hampshire; JBG – Jersey Black Giant; RIR – Rodhe Island Red; IG – Indigenous Giant; IG × NHS – F1 generation resulted from the cross between Indigenous Giant and New Hampshire; IG × JBG – F1 generation resulted from the cross between Indigenous Giant and Jersey Black Giant. ⁽¹⁾Edible viscera; ⁽²⁾Abdominal fat.

Regarding the weight of edible viscera, an interaction was observed between genotype and sex (Table 3), with males of all genotypes showing higher values (Table 7). In fact, males present a greater food digestion efficiency since they have a higher body weight, making the organs involved in this process, such as liver and gizzard, to be more developed (MITROVIC et al., 2011). For edible viscera yield, a significant effect was observed only for genotype (Table 3), with the genotype RIR presenting higher yield when compared to other breeds (Table 7). This is probably due to their lower average carcass weight (Table 4), which may have increased the proportion and, consequently, the yield of edible viscera.

An interaction was observed between genotype and sex for abdominal fat yield (Table 3). In this sense, a difference between sexes was observed only for NHS and JBG, with females presenting higher values (Table 7). Santos et al. (2005), Faria et al. (2011), Mitrovic et al. (2011), and Sarica et al. (2014) observed a higher average of abdominal fat yield in females, which is probably due to differences in growth rates between sexes. Because females present earlier maturity, they deposit a greater amount of fat in the carcass, mainly in the abdominal region (GONZALES; SARTORI, 2002).

For both sexes, the genotypes NHS and JBG presented higher abdominal fat yields (Table 7). An effect of genotype was observed on abdominal fat weight (Table 3), with the genotypes NHS and JBG showing the highest averages (Table 7). The fastest-growing broiler chickens achieved a peak of muscle growth and physiological maturity earlier than slow-growing broiler chickens. As a result, a growing portion of diet energy is converted into adipose tissue, which will accumulate first in the abdominal region (GONZALES; SARTORI, 2002). Thus, the fastest-growing genotypes, such as NHS and JBG, tend to have a higher proportion of abdominal fat due to the lipid deposition be earlier in these broiler chickens.

Conclusions

The genotypes resulting from the crossing of different breeds of broiler chickens provided better results for meat production, with a higher growth potential, carcass characteristics, and muscular portion of both cuts.

In the production of chicken meat under an alternative system, females demonstrated a higher breast yield and a higher amount of meat in the leg despite producing lighter carcasses.

References

- ALBINO, L. F. T.; TAVERNARI, F. C.; VIEIRA, R. A.; NERY, L. R. Criação de frango e galinha caipira, sistema alternativo de criação de aves. In: ALBINO, L. F. T.; TAVERNARI, F. C.; VIEIRA, R. A.; NERY, L. R. (Ed.). *Aves mais indicadas*. Viçosa: Aprenda Fácil, 2014. p. 21-38.
- ASSOCIAÇÃO BRASILEIRA DOS CRIADORES DE INDIO GIGANTE – ABCIG. Manual do criador de Índio Gigante. Belém: Ancrib, 2016. Disponível em: <http://abcig.com.br/manual_do_criador_indio_gigante>. Acesso em: 15 maio 2016.
- DEL CASTILHO, C. C.; SANTOS, T. T.; RODRIGUES, C. A. F.; TORRES FILHO, R. A. Effects of sex and genotype on performance and yield characteristics of free range broiler chickens. *Arquivo Brasileiro de Medicina Veterinária e Zootecnia*, Belo Horizonte, v. 65, n. 5, p. 1483-1490, 2013. Disponível em: <<http://www.scielo.br/pdf/abmvz/v65n5/a29v65n5.pdf>>. Acesso em: 23 fev. 2016.
- FARIA, P. B.; BRESSAN, M. C.; SOUZA, X. R.; ROSSATO, L. V.; BOTEGA L. M. G.; GAMA L. T. Carcass and parts yield of broilers reared under a semi-extensive system. *Revista Brasileira de Ciência Avícola / Brazilian Journal of Poultry Science*, Campinas, v. 12, n. 3, p. 153-159, 2010. Disponível em: <<http://www.scielo.br/pdf/rbca/v12n3/03.pdf>>. Acesso em: 21 mar. 2016.
- FARIA, P. B.; VIEIRA, J. O.; SILVA, J. N.; RODRIGUES, A. Q.; SOUZA, X. R.; SANTOS, F. R.; PEREIRA, A. A. Performance and carcass characteristics of alternative broiler chickens fed diets containing alternative foods. *Revista Brasileira de Ciência Avícola / Brazilian Journal of Poultry Science*, Campinas, v. 13, n. 3, p. 113-118, 2011. Disponível em: <<http://www.scielo.br/pdf/rbca/v13n3/09.pdf>>. Acesso em: 21 mar. 2016.

- GONZALES, E.; BUYSE, J.; TAKITA, T. S.; SARTORI, J. R.; DECUYPERE, E. Metabolic disturbances in male broilers of different strains. 1. Performance, mortality and right ventricular hypertrophy. *Poultry Science*, Champaign, v. 77, n. 11, p. 1646-1653, 1998. Disponível em: <<https://academic.oup.com/ps/article-lookup/doi/10.1093/ps/77.11.1646>>. Acesso em: 25 jun. 2016.
- GONZALES, E.; SARTORI, J. R. Crescimento e metabolismo muscular. In: MACARI, M.; FURLAN, R. L.; GONZALES, E. (Ed.). *Fisiologia aviária aplicada a frangos de corte*. Jaboticabal: FUNEP/UNESP, 2002. p. 279-297.
- ISOLA, J. G. M. P.; MORAES, P. C.; RAHAL, S. C.; MACHADO, M. R. F. Morfologia, ultraestrutura e morfometria do tegumento da paca (*Cuniculus paca* Linnaeus, 1766) criada em cativeiro. *Pesquisa Veterinária Brasileira*, Seropédica, v. 33, n. 5, p. 674-682, 2013. Disponível em: <<http://repositorio.unesp.br/bitstream/handle/11449/75290/2-s2.0-84880783928.sequence=1&isAllowed=y>>. Acesso em: 20 maio 2016.
- KESSLER, A. M.; SNIZEK, P. N.; BRUGALLI, I. Manipulação da quantidade de gordura na carcaça de frangos. In: FUNDAÇÃO DE CIÊNCIA E TECNOLOGIA AVÍCOLAS, 1., 2000, Campinas. *Anais...* Campinas: Fundação Apinco de Ciência e Tecnologia Avícolas, Campinas, 2000. p. 107-133.
- MADEIRA, L. A.; SARTORI, J. R.; ARAUJO, P. C.; PIZZOLANTE, C. C.; SALDANHA, E. S. P. B.; PEZZATO, A. C. Morfologia das fibras musculares esqueléticas de frangos de corte de diferentes linhagens criados em sistema de confinamento e semi confinamento. *Revista Brasileira de Zootecnia*, Viçosa, v. 35, n. 6, p. 2322-2332, 2006. Disponível em: <<http://www.scielo.br/pdf/0D/rbz/v35n6/18.pdf>>. Acesso em: 13 abr. 2016.
- MIKULSKI, D.; CELEJ, J.; JANKOWSKI, J.; MAJEWSKA, T.; MIKULSKAET, M. Growth performance, carcass traits and meat quality of slower-growing and fast-growing chickens raised with and without outdoor access. *Asian-Australasian Journal Animal Science*, Seoul, v. 24, n. 10, p. 1407-1416, 2011. Disponível em: <http://ocean.kisti.re.kr/download/volume/aaaaps/E1DMBP/2011/v24n10/E1DMBP_2011_v24n10_1407.pdf>. Acesso em: 14 abr. 2016.
- MITROVIC, S.; BOGOSAVLJEVIC-BOSKOVIC, S.; STANISIC, G.; DJERMANOVIC, V.; DOSKOVIC, V.; RAKONJAC, S. Carcass characteristics of two strains of native broilers (White Naked Neck and Black Svrljig) fattened under a semi-intensive system. *African Journal of Biotechnology*, Nairobi, v. 10, n. 70, p. 15813-15818, 2011. Disponível em: <<http://search.proquest.com/openview/ba1d6bef9420f92a212ba2d7ce5dca30/1?pq-origsite=gscholar&cbl=1006374>>. Acesso em: 14 abr. 2016.
- SANTOS, A. L.; SAKOMURA, N. K.; FREITAS, E. R.; FORTES, C. M. L.; CARRILHO, E. N. V. M.; FERNANDES, J. B. K. Estudo do crescimento, desempenho, rendimento de carcaça e qualidade de carne de três linhagens de frango de corte. *Revista Brasileira de Zootecnia*, Viçosa, v. 34, n. 5, p. 1589-1598, 2005. Disponível em: <<http://www.scielo.br/pdf/rbz/v34n5/26640.pdf>>. Acesso em: 25 abr. 2016.
- SARICA, M.; YAMAK, U. S.; TURHAN, S.; BOZ, M. A.; SARICAOGLU, F. T.; ALTOP, A. Comparing slow-growing chickens produced by two- and three-way crossings with commercial genotypes. 2. Carcass quality and blood parameters. *European Poultry Science*, Stuttgart, v. 14, n. 1, p. 2014-2030, 2014. Disponível em: <https://www.researchgate.net/profile/Furkan_Turker/publication/262635967_Comparing_slowgrowing_chickens_produced_by_twoand_threeway_crossings_with_commercial_genotypes_2_Carcass_quality_and_blood_parameters/links/0deec53848776dc1eb000000/Comparing-slow-growing-chickens-produced-by-two-and-three-way-crossings-with-commercial-genotypes-2-Carcass-quality-and-blood-parameters.pdf>. Acesso em: 15 abr. 2016.
- SOGUNLE, O. M.; OLANIYI, O. A.; EGBEYALE, L. T.; AKINOLA, O. S.; SHITTU, T. A.; ABIOLA, S. S.; LADOKUN, A. O.; SOBAYO, R. A. Free range and deep litter poultry production systems: effect on performance, carcass yield and meat composition of cockerel chickens. *Tropical Animal Health and Production*, Edinburgh, v. 45, n. 1, p. 281-288, 2013. Disponível em: <<https://link.springer.com/article/10.1007/s11250-012-0213-2>>. Acesso em: 16 abr. 2016.
- VELOSO, R. C.; PIRES, A. V.; TORRES FILHO, R. A.; PINHEIRO, S. R. F.; WINKELSTROTER, L. K.; ALCÂNTARA, D. C.; CRUZ, C. C. D. C. S. Parâmetros de desempenho e carcaça de genótipos de frangos tipo caipira. *Arquivo Brasileiro de Medicina Veterinária e Zootecnia*, Belo Horizonte, v. 66, n. 4, p. 1251-1259, 2014. Disponível em: <<http://www.scielo.br/pdf/abmvz/v66n4/0102-0935-abmvz-66-04-01251.pdf>>. Acesso em: 02 maio 2016.