

## Sample size to estimate the mean of morphological traits of rye cultivars in sowing dates and evaluation times

### Tamanho de amostra para estimação da média de caracteres morfológicos de cultivares de centeio em épocas de semeadura e de avaliação

Cirineu Tolfo Bandeira<sup>1</sup>; Alberto Cargnelutti Filho<sup>2\*</sup>; Diego Nicolau Follmann<sup>2</sup>; Cláudia Marques de Bem<sup>3</sup>; Cleiton Antonio Wartha<sup>4</sup>; Rosana Marzari Thomasi<sup>5</sup>

#### Abstract

The objectives of this study were to determine the sample size (number of plants) required to estimate the mean of morphological traits of rye (*Secale cereale* L.) and verify the sample size variability between the traits, cultivars, sowing dates, and evaluation times for distinct mean estimation errors. Ten uniformity trials were performed with two rye cultivars (BRS Progresso and Temprano) in five sowing dates (05/03/2016, 05/25/2016, 06/07/2016, 06/22/2016 and 07/04/2016). Evaluations of traits plant height, number of leaves, and number of stems were performed during the development of the crop. In order to verify the difference of the traits between cultivars and between sowing dates and evaluation times, the F-test was applied to test the hypothesis of homogeneity of variances and the Student's t-test was used to test the hypothesis of equality of means. The sample size of each trait was calculated for distinct mean estimation errors. There is sample size variability between the traits, cultivars, sowing dates, and evaluation times. In order to estimate the mean of plant height with the same precision, smaller sample sizes are required at the intermediate and final evaluation times compared to initial evaluation times. For the traits number of leaves and stems, smaller sample sizes are required in the initial evaluation times than in the final evaluation times. For mean estimation of traits with maximum estimation error of 15% between sowing dates and evaluation times, 83 and 103 plants are required respectively for cultivars BRS Progresso and Temprano.

**Key words:** Sampling planning. Sampling precision. *Secale cereale* L.

#### Resumo

Os objetivos deste trabalho foram determinar o tamanho de amostra (número de plantas) para a estimação da média de caracteres morfológicos de centeio (*Secale cereale* L.) e verificar a variabilidade do tamanho de amostra entre caracteres, entre cultivares, entre épocas de semeadura e de avaliação, para distintos erros de estimação da média. Foram conduzidos dez ensaios de uniformidade, com duas cultivares de centeio (BRS Progresso e Temprano) em cinco épocas de semeadura (03/05/2016,

<sup>1</sup> Eng<sup>o</sup> Agr<sup>o</sup>, Discente de Mestrado, Programa de Pós-Graduação em Agronomia, Universidade Federal de Santa Maria, UFSM, Santa Maria, RS, Brasil. E-mail: cirineutolfobandeira@gmail.com

<sup>2</sup> Eng<sup>os</sup> Agr<sup>os</sup>, Profs. Drs., Departamento de Fitotecnia, UFSM, Santa Maria, RS, Brasil. E-mail: alberto.cargnelutti.filho@gmail.com; diegonicolaufollmann@gmail.com

<sup>3</sup> Zootecnista, Dr<sup>a</sup>, Programa de Pós-Graduação em Agronomia, UFSM, Santa Maria, RS, Brasil. E-mail: claudia\_debem@hotmail.com

<sup>4</sup> Eng<sup>o</sup> Agr<sup>o</sup>, Discente de Mestrado, Programa de Pós-Graduação em Fitotecnia, Universidade Federal de Viçosa, UFV, Viçosa, MG, Brasil. E-mail: cleiton.ufsm@gmail.com

<sup>5</sup> Discente de Graduação em Agronomia, UFSM, Santa Maria, RS, Brasil. E-mail: rosanamthomasi@hotmail.com

\* Author for correspondence

25/05/2016, 07/06/2016, 22/06/2016 e 04/07/2016). Durante o desenvolvimento da cultura foram realizadas avaliações dos caracteres altura de planta, número de folhas e número de colmos. Para verificar as diferenças dos caracteres, entre cultivares e entre épocas de semeadura e de avaliação, foi aplicado o teste F, para testar a hipótese de homogeneidade de variâncias e o teste t de Student, para testar a hipótese de igualdade de médias. Foi calculado o tamanho de amostra de cada caractere para distintos erros de estimação da média. Há variabilidade do tamanho de amostra entre os caracteres, cultivares, épocas de semeadura e épocas de avaliação. Para a estimação da média de altura de planta, com mesma precisão, é necessário menor tamanho de amostra nas épocas de avaliação intermediárias e finais comparadas as épocas iniciais de avaliação. Para os caracteres números de folhas e de colmos nas épocas iniciais de avaliação, é necessário tamanho de amostra menor do que em épocas finais de avaliação. São necessárias 83 e 103 plantas, respectivamente, para as cultivares BRS Progresso e Temprano, para a estimação da média dos caracteres, com erro máximo de estimação de 15%, entre épocas de semeadura e entre épocas de avaliação.

**Palavras-chave:** Planejamento amostral. Precisão amostral. *Secale cereale* L.

## Introduction

Rye (*Secale cereale* L.) belongs to the Poaceae family and is planted annually during winter and matures earlier than many other cereals, but has a long reproductive cycle (BAIER, 1994). In Brazil, the primary states that grow rye are Rio Grande do Sul and Paraná; the cultivation area of rye is approximately 2,500 ha and average yield of rye is 2,600 kg ha<sup>-1</sup> (CONAB, 2017). Rye has vigorous initial growth and has high productivity as measured in dry matter weight. In the agricultural sector, rye is important as a forage crop, cover crop, and is an important production grain. Its straw has a high carbon: nitrogen ratio and rye is suitable as a soil cover crop, as well as in the management of soil erosion and reduction of NO<sub>3</sub> losses (PANTOJA et al., 2016). The use of grasses in monoculture or in consortium has a greater potential for soil protection because of the prolonged presence of their remnants on the soil surface (ZIECH et al., 2015).

The high demands of time, labor, and financial resources hamper the evaluation and measurement of a large number of traits of plants in agricultural experiments. Thus, efficient sampling for the desired measurements is needed. Further, sample size is important in obtaining accurate estimates of the parameters in question. The sample size needed depends on the variability in the data and the desired degree of confidence, which are determined by the

researcher (BUSSAB; MORETTIN, 2013). Sample size interferes with the interpretation of significance of statistical tests, as small samples can generate inaccurate estimates, which are less likely to occur with large sample sizes. However, larger samples can generate deflated *P*-values (i.e., indicating statistical significance in the absence of practical significance (LIN et al., 2013).

Scenarios formed by cultivars, sowing dates, and evaluation times during the crop cycle are important for the determination of sample size. Data that cover the range of variation in traits of interest better represent the crop, and these data help generate the capacity to estimate needed sample sizes that can be used as a reference for planning of future experiments. The need for variability in sample sizes to estimate means and associated errors for plant traits, cultivars, sowing dates, and evaluation times have been reported for crops such as maize (STORCK et al., 2007), castor bean (CARGNELUTTI FILHO et al., 2010), black oat (CARGNELUTTI FILHO et al., 2015), pigeon pea (FACCO et al., 2015, 2016), *Crotalaria juncea*, and *Crotalaria spectabilis* Roth (TEODORO et al., 2015).

Studies regarding sample sizes needed to estimate the mean of morphological traits in rye for BRS Progresso and Temprano cultivars were not found in the literature and we assumed that

there is variability in sample size needed to assess traits, cultivars, sowing dates, and evaluation times. The objectives of this study were to determine the sample size (number of plants) required to estimate the means of morphological traits of rye (*Secale cereale* L.) and verify that for a given level of statistical confidence, sample sizes varies for traits, cultivars, sowing dates, and evaluation times.

## Material and Methods

Ten uniformity trials (blank experiments) were performed with rye (*Secale cereale* L.), in the experimental area of the Department of Plant Science at the Federal University of Santa Maria, Rio Grande do Sul in Southern Brazil (29°42'S, 53°49'W; elevation = 95 m) during the 2016 growing season. According to the Köppen Climate Classification, the climate of the region is Cfa subtropical humid, with hot summers and no defined dry season (HELDWEIN et al., 2009). The soil is classified as sandy loam typic Paleudalf (SANTOS et al., 2013).

Conventional soil tillage was carried out with light harrowing and basic fertilization was performed with 500 kg ha<sup>-1</sup> of NPK fertilizer (5-20-20) being applied. In these uniformity trials, procedures of sowing, fertilization, cultural practices, and evaluations were performed homogeneously throughout the experimental area. Two cultivars (BRS Progresso and Temprano) were sown at five dates: 05/03/2016 (Sowing date 1), 05/25/2016 (Sowing date 2), 06/07/2016 (Sowing date 3), 06/22/2016 (Sowing date 4), and 04/07/2016 (Sowing date 5). Seeds from the BRS Progresso and Temprano cultivars were sown by broadcasting at a seed density of 455 seeds m<sup>-2</sup>. The choice of cultivars was based on their suitability for use in this region. The BRS Progresso cultivar is destined for grain production and Temprano is recommended as a forage and cover crop. Each cultivar was sown in an area of 320 m<sup>2</sup> (20 m × 16 m) at the first sowing

date, whereas each cultivar occupied 375 m<sup>2</sup> (25 m × 15 m) at the other sowing dates.

In the central area of 100 m<sup>2</sup> (10 m × 10 m) in each uniformity trial, a grid with 100 sample points was marked with stakes spaced at 1 m × 1 m, forming a matrix of ten rows and ten columns. The closest plant to the sampling stake was measured at each sampling point. In these sampled plants, plant height (PH, in cm; measured as the distance from the soil surface to the insertion of the last leaf on the stem), number of leaves per plant (NL; leaves of the main stem + leaves of the tillers), and the number of stems per plant (NS; main stem + stems of the tillers) were recorded.

Evaluations were carried out from the beginning of crop development until flowering (Table 1). Measurements were collected using the same methodology for all sowing dates, respecting the dates of planting and plant development for each sowing date.

The mean ( $\bar{m}$ ) and variance ( $s^2$ ) were calculated at each sowing date for each trait, cultivar, and evaluation time. The  $F$ -test (unilateral), at 5% of error probability was used to test homogeneity of variance and the  $t$ -test (bilateral) at 5% of error probability for independent samples was performed to test the hypothesis of equality of means. Initially, the tests were applied to the data from the last evaluation (i.e., the evaluation performed at the flowering stage of the crop). The cultivars were compared for each sowing date with these data. Afterwards, the sowing dates were compared in each cultivar. Then, with data from all evaluations, evaluation times were compared within each combination of cultivar and sowing date. Non-normally distributed data were assessed using the Kolmogorov-Smirnov test and data randomness was checked by the run test (CAMPOS, 1983). The sample size ( $n$ ) was calculated for the confidence interval half-lengths (estimation errors) of 5, 10, 15, 20, and 25% of the estimate of the mean with confidence level ( $1-\alpha$ ) of 95%, following

the formula:  $n = \frac{t_{\alpha/2}^2 s^2}{(\text{estimation error})^2}$  (BUSSAB; MORETTIN, 2013), where:  $t_{\alpha/2}$  is the critical value of the Student's  $t$  distribution, whose area on the right

is equal to  $\alpha/2$  (i.e., the value of  $t$  such that:  $P(t > t_{\alpha/2}) = \alpha/2$ , with  $(n-1)$  degrees of freedom, with  $\alpha = 5\%$  of probability of error) and,  $s^2$  is the estimate of variance.

**Table 1.** Evaluation times in days after sowing of the traits plant height, number of leaves, and number of stems in two rye cultivars in five sowing dates.

Evaluation	Sowing date				
	1 (05/03/2016)	2 (05/25/2016)	3 (06/07/2016)	4 (06/22/2016)	5 (07/04/2016)
-----BRS Progresso-----					
1	22	20	22	16	24
2	29	28	28	20	29
3	36	34	35	27	36
4	42	41	42	36	43
5	50	49	51	41	50
6	56	55	56	50	58
7	66	65	63	55	66
8	71	69	73	64	72
9	97	99	100	92	87
-----Temprano-----					
1	22	20	22	16	24
2	29	28	28	20	29
3	36	34	35	27	36
4	42	41	42	36	43
5	50	49	51	41	50
6	56	55	56	50	58
7	66	65	63	55	66
8	71	69	72	63	72
9	92	92	93	85	95
10	113	131	125	121	120
11	142	-	-	-	-

Based on the formula used to calculate the sample size,  $n$  was set at 100 plants and the estimation error was calculated as a percentage of the estimate of the mean ( $m$ ) for each trait, using the formula:  $\text{estimation error} = 100 \frac{t_{\alpha/2} s}{\sqrt{n} m}$ , where  $s$  is the estimate of standard deviation. Statistical analyses were performed using GENES software (CRUZ, 2013) and Microsoft Excel®.

## Results and Discussion

There were differences between the means of plant height for the cultivars at the flowering stage shown by the Student's  $t$ -test only for the first sowing date. The number of leaves per plant was greater in the BRS Progresso cultivar for sowing dates 1 and 4. However, there was superiority of the Temprano cultivar over BRS Progresso for the number of stems

per plant at sowing dates 2, 4, and 5. The variances were heterogeneous for 80% of the cases, indicating that the sample size should be different between the cultivars (Table 2). Differences in sample size were observed between maize (STORCK et al.,

2007) and soybean genotypes (CARGNELUTTI FILHO et al., 2009), between castor bean hybrids (CARGNELUTTI FILHO et al., 2010), and between species of crotalaria (TEODORO et al., 2015).

**Table 2.** Mean and variance of traits plant height (PH, in cm), number of leaves per plant (NL), and number of stems per plant (NS) at the flowering stage of two rye cultivars (BRS Progresso and Temprano), Student's t-test results for comparison of means, and F-test for comparison of variances of rye cultivars in five sowing dates.

Trait	-----Mean-----		t-test	-----Variance-----		
	BRS Progresso	Temprano		BRS Progresso	Temprano	F-test
Sowing date 1 (05/03/2016)						
PH	101.83	114.32	*	158.15	294.01	Heterogeneous
NL	10.98	7.74	*	41.43	20.76	Heterogeneous
NS	2.81	2.72	ns	2.58	2.51	Homogeneous
Sowing date 2 (05/25/2016)						
PH	118.60	120.97	ns	299.70	480.88	Heterogeneous
NL	20.30	20.97	ns	110.01	189.77	Heterogeneous
NS	5.08	6.33	*	5.31	23.40	Heterogeneous
Sowing date 3 (06/07/2016)						
PH	126.44	126.91	ns	183.99	434.83	Heterogeneous
NL	11.62	12.64	ns	25.73	49.30	Heterogeneous
NS	3.28	3.67	ns	2.37	3.64	Heterogeneous
Sowing date 4 (06/22/2016)						
PH	118.34	122.56	ns	158.09	469.35	Heterogeneous
NL	10.73	7.65	*	23.69	18.80	Homogeneous
NS	2.83	3.71	*	1.27	5.22	Heterogeneous
Sowing date 5 (07/04/2016)						
PH	107.92	110.52	ns	340.21	356.70	Homogeneous
NL	6.68	7.49	ns	11.49	31.55	Heterogeneous
NS	2.08	3.32	*	1.35	4.06	Heterogeneous

\*Means differ by bilateral t-test at 5% of error probability. ns: not significant. Heterogeneous: heterogeneous variances by bilateral F-test at 5% probability of error. Homogeneous: homogeneous variances.

There was a distinct behavior of cultivars between sowing dates. The difference between mean PH values was not significant in one case (sowing dates 2 and 4) and in three cases (sowing dates 1 and 5, 2 and 4, and 3 and 4) respectively for BRS Progresso and Temprano (Table3). Therefore, the BRS Progresso cultivar was more influenced by the environment than Temprano for the PH

trait. The NL mean in the BRS Progresso cultivar was different for many sowing dates, except in the comparison of sowing date 1 with sowing dates 3 and 4, and sowing date 3 with sowing date 4. Meanwhile, Temprano means were not different between sowing dates 1 and 4, or 1 and 5, and sowing dates 4 and 5. For the NS trait, the mean was not different between sowing dates 1 and 4 for

BRS Progresso. The same trait was not different for Temprano between sowing dates 3 and 4, 3 and 5, and 4 and 5. This demonstrated that sowing date affected the traits in question, as the lowest values were found consistently for the first and last

sowing dates. When comparing variances, at least in 60% and 30% of the comparisons respectively for BRS Progresso and Temprano, variances were heterogeneous, which suggests different sample sizes are needed for different sowing dates.

**Table 3.** Results of Student's t test (above diagonal) for comparison of means (diagonal) and F-test (below diagonal) for comparison of variances between sowing dates in rye cultivars BRS Progresso and Temprano at flowering stage.

	----- BRS Progresso -----						----- Temprano -----				
	1	2	3	4	5		1	2	3	4	5
Plant height											
1	101.83 <sup>(1)</sup>	*	*	*	*	114.32	*	*	*		ns
2	Het	118.60	*	ns	*	Het	120.97	*		ns	*
3	Hom	Het	126.44	*	*	Het	Hom	126.91		ns	*
4	Hom	Het	Hom	118.34	*	Het	Hom	Hom	122.56		*
5	Het	Hom	Het	Het	107.92	Hom	Hom	Hom	Hom	Hom	110.52
Number of leaves per plant											
1	10.98	*	ns	ns	*	7.74	*	*		ns	ns
2	Het	20.30	*	*	*	Het	20.97	*	*	*	*
3	Het	Het	11.62	ns	*	Het	Het	12.64	*	*	*
4	Het	Het	Hom	10.73	*	Hom	Het	Het	7.65		ns
5	Het	Het	Het	Het	6.68	Het	Het	Het	Het	Het	7.49
Number of stems per plant											
1	2.81	*	*	ns	*	2.72	*	*	*	*	*
2	Het	5.08	*	*	*	Het	6.33	*	*	*	*
3	Hom	Het	3.28	*	*	Het	Het	3.67		ns	ns
4	Het	Het	Het	2.83	*	Het	Het	Het	3.71		ns
5	Het	Het	Het	Hom	2.08	Het	Het	Hom	Hom	Hom	3.32

\*Means differ by bilateral t-test at 5% of error probability. ns: not significant. Heterogeneous: heterogeneous variances by bilateral F-test at 5% probability of error. Homogeneous: homogeneous variances. <sup>(1)</sup>Mean of traits of the rye cultivars in sowing dates at flowering stage.

There were differences for PH in both cultivars and for all sowing dates, showing that there was a difference between evaluation times for each cultivar. There were differences between NL and NS traits in BRS Progresso and Temprano cultivars for most evaluation times. When comparing the variances at evaluation times for BRS Progresso, the variances were heterogeneous at least in 88.89, 80.56, and 63.89% of the comparisons respectively for PH, NL, and NS traits. For the Temprano cultivar,

the variances were heterogeneous at least in 89.09, 88.89, and 68.89% of the comparisons respectively for PH, NL, and NS traits (Table 4). Therefore, the sample size to estimate the mean should be different between the evaluation times.

These results corroborate those reported by Facco et al. (2015), who verified that the optimal sample size in pigeon pea depends on the morphological trait being investigated and evaluation time. Also,

the needed sample size varied with the productive trait and harvest for this same crop (FACCO et al., 2016).

The data adjusted to the normal distribution by the Kolmogorov-Smirnov test and were randomized according to the run test (CAMPOS,

1983) respectively in 53.13% and 76.39% of the 288 cases (3 traits × 2 cultivars × 5 sowing dates × 9, 10, or 11 evaluation times). The sample size was calculated based on the Student's *t* distribution as the majority of cases had normally distributed data and randomness associated with the great number of plants sampled in each case (*n* = 100 plants).

**Table 4.** Number of significant (\*) and non-significant (ns) results by the Student's t-test and number of results of comparison of variances (heterogeneous and homogeneous) by the F-test during the evaluation times of plant height (PH), number of leaves per plant (NL), and number of stems per plant (NS) of rye cultivars BRS Progresso and Temprano.

Trait	----- BRS Progresso <sup>(1)</sup> -----				----- Temprano <sup>(2)</sup> -----			
	-- t-test --		----- F-test -----		-- t-test --		----- F-test -----	
	*	ns	Het	Hom	*	ns	Het	Hom
Sowing date 1 (05/03/2016)								
PH	36	0	34	2	55	0	49	6
NL	34	2	33	3	53	2	51	4
NS	28	8	31	5	52	3	51	4
Sowing date 2 (05/25/2016)								
PH	36	0	36	0	45	0	43	2
NL	32	4	31	5	41	4	45	0
NS	31	5	24	12	42	3	31	14
Sowing date 3 (06/07/2016)								
PH	36	0	35	1	45	0	43	2
NL	31	5	29	7	39	6	40	5
NS	33	3	23	13	41	4	32	13
Sowing date 4 (06/22/2016)								
PH	36	0	32	4	45	0	41	4
NL	32	4	34	2	43	2	40	5
NS	29	7	25	11	43	2	33	12
Sowing date 5 (07/04/2016)								
PH	36	0	33	3	45	0	45	0
NL	29	7	30	6	42	3	43	2
NS	31	5	26	10	38	7	37	8

\*Means differ by bilateral t-test at 5% of error probability. ns: not significant. Heterogeneous: heterogeneous variances by bilateral F-test at 5% probability of error. Homogeneous: homogeneous variances. (1) In this cultivar, 36 combinations of evaluation times are possible in each sowing date for comparison of means. (2) In this cultivar, 55 and 45 combinations of evaluation times are possible respectively for sowing date 1 and other sowing dates for comparison of means.

The sample size for the estimation of the mean of each trait in evaluation times, based on a 95% confidence interval, ranged from 18 plants for PH traits in the ninth evaluation (flowering) of sowing date 4 in the BRS Progresso cultivar to 920 plants for NS trait in the tenth evaluation of sowing date 2 in the Temprano cultivar (Tables 5, 6). Therefore, there was great variability in sample size between traits, cultivars, sowing dates, and evaluation times. Thus, 920 rye plants are required to assess differences in means of the traits, cultivars, sowing dates, or evaluation times, at a confidence level of 95% and an error of 5% of the estimate of the mean.

From a practical point of view, there are difficulties in evaluating 920 rye plants. Thus, sample sizes with 10, 15, 20, and 25% estimation errors were determined (Tables 5, 6). Accepting larger estimation errors provides researchers the flexibility to use sample sizes according to the desired accuracy to meet research needs. Based on the largest sample size between the traits PH, NL, and NS, cultivars, sowing dates, and evaluation times with an estimation error of 15% and 95% level of confidence, 103 plants are required to evaluate the mean of these traits. Thus, in an experimental design with four replications, 26 plants per replication of each treatment should be evaluated.

**Table 5.** Sample size (number of plants) to estimate the mean of the traits plant height, number of leaves per plant, and number of stems per plant in the rye cultivar BRS Progresso for the estimation errors equal to 5% , 10%, 15%, 20%, and 25% of the mean (m) in sowing dates and evaluation times (E).

E	Plant height						Number of leaves per plant						Number of stems per plant					
	5	10	15	20	25	Error (%)	5	10	15	20	25	Error (%)	5	10	15	20	25	Error (%)
Sowing date 1 (05/03/2016)																		
1	220	55	25	14	9	7.41	92	23	11	6	4	4.77	196	49	22	13	8	6.99
2	123	31	14	8	5	5.52	94	24	11	6	4	4.84	185	47	21	12	8	6.79
3	109	28	13	7	5	5.22	105	27	12	7	5	5.12	203	51	23	13	9	7.12
4	136	34	16	9	6	5.82	175	44	20	11	7	6.61	194	49	22	13	8	6.96
5	165	42	19	11	7	6.41	226	57	26	15	10	7.50	253	64	29	16	11	7.94
6	180	45	20	12	8	6.70	199	50	23	13	8	7.05	448	112	50	28	18	10.58
7	189	48	21	12	8	6.86	428	107	48	27	18	10.34	440	110	49	28	18	10.48
8	149	38	17	10	6	6.09	290	73	33	19	12	8.51	427	107	48	27	18	10.33
9	25	7	3	2	1	2.45	542	136	61	34	22	11.63	515	129	58	33	21	11.34
Sowing date 2 (05/25/2016)																		
1	159	40	18	10	7	6.29	44	11	5	3	2	3.30	-( <sup>1</sup> )	-	-	-	-	-
2	116	29	13	8	5	5.37	84	21	10	6	4	4.57	49	13	6	4	2	3.47
3	117	30	13	8	5	5.39	96	24	11	6	4	4.90	124	31	14	8	5	5.57
4	96	24	11	6	4	4.88	143	36	16	9	6	5.98	175	44	20	11	7	6.60
5	111	28	13	7	5	5.27	161	41	18	11	7	6.33	323	81	36	21	13	8.97
6	120	30	14	8	5	5.48	216	54	24	14	9	7.34	357	90	40	23	15	9.44
7	131	33	15	9	6	5.70	320	80	36	20	13	8.94	742	186	83	47	30	13.61
8	120	30	14	8	5	5.48	297	75	33	19	12	8.61	420	105	47	27	17	10.24
9	34	9	4	3	2	2.90	421	106	47	27	17	10.25	324	81	36	21	13	9.00

continue

continuation

Sowing date 3 (06/07/2016)																		
1	131	33	15	9	6	5.72	188	47	21	12	8	6.84	-	-	-	-	-	-
2	82	21	10	6	4	4.51	48	12	6	3	2	3.45	76	19	9	5	4	4.34
3	75	19	9	5	3	4.32	71	18	8	5	3	4.20	60	15	7	4	3	3.86
4	92	23	11	6	4	4.78	74	19	9	5	3	4.30	102	26	12	7	5	5.03
5	68	17	8	5	3	4.10	142	36	16	9	6	5.96	237	60	27	15	10	7.69
6	66	17	8	5	3	4.05	215	54	24	14	9	7.31	182	46	21	12	8	6.74
7	88	22	10	6	4	4.68	204	51	23	13	9	7.14	196	49	22	13	8	7.00
8	43	11	5	3	2	3.27	196	49	22	13	8	7.00	203	51	23	13	9	7.12
9	19	5	3	2	1	2.13	301	76	34	19	13	8.66	347	87	39	22	14	9.30
Sowing date 4 (06/22/2016)																		
1	149	38	17	10	6	6.08	127	32	15	8	6	5.61	-	-	-	-	-	-
2	70	18	8	5	3	4.17	50	13	6	4	2	3.51	107	27	12	7	5	5.15
3	48	12	6	3	2	3.44	89	23	10	6	4	4.71	121	31	14	8	5	5.48
4	89	23	10	6	4	4.71	90	23	10	6	4	4.72	123	31	14	8	5	5.53
5	83	21	10	6	4	4.54	106	27	12	7	5	5.13	139	35	16	9	6	5.88
6	87	22	10	6	4	4.64	105	27	12	7	5	5.10	166	42	19	11	7	6.44
7	118	30	14	8	5	5.41	159	40	18	10	7	6.30	168	42	19	11	7	6.47
8	69	18	8	5	3	4.14	221	56	25	14	9	7.43	289	73	33	19	12	8.49
9	18	5	2	2	1	2.11	325	82	37	21	13	9.00	251	63	28	16	11	7.91
Sowing date 5 (07/04/2016)																		
1	92	23	11	6	4	4.79	58	15	7	4	3	3.78	172	43	20	11	7	6.55
2	108	27	12	7	5	5.19	46	12	6	3	2	3.37	65	17	8	5	3	4.00
3	83	21	10	6	4	4.55	112	28	13	7	5	5.27	89	23	10	6	4	4.71
4	107	27	12	7	5	5.17	173	44	20	11	7	6.57	203	51	23	13	9	7.12
5	140	35	16	9	6	5.92	235	59	27	15	10	7.66	203	51	23	13	9	7.12
6	114	29	13	8	5	5.32	249	63	28	16	10	7.88	291	73	33	19	12	8.53
7	81	21	9	6	4	4.49	280	70	32	18	12	8.35	381	96	43	24	16	9.75
8	65	17	8	5	3	4.02	416	104	47	26	17	10.19	415	104	47	26	17	10.18
9	47	12	6	3	2	3.39	406	102	46	26	17	10.07	491	123	55	31	20	11.07

(1) It was not possible to calculate the sample size because there was no variance in the data (plants presented only one stem).

**Table 6.** Sample size (number of plants) to estimate the mean of the traits plant height, number of leaves per plant, and number of stems per plant in the rye cultivar Temprano for the estimation errors equal to 5%, 10%, 15%, 20%, and 25% of the mean (m) in sowing dates and evaluation times (E).

E	Plant height						Number of leaves per plant						Number of stems per plant					
	5	10	15	20	25	Error (%)	5	10	15	20	25	Error (%)	5	10	15	20	25	Error (%)
Sowing date 1 (05/03/2016)																		
1	111	28	13	7	5	5.26	102	26	12	7	5	5.03	160	40	18	10	7	6.31
2	78	20	9	5	4	4.39	87	22	10	6	4	4.66	135	34	15	9	6	5.80
3	70	18	8	5	3	4.17	80	20	9	5	4	4.46	81	21	9	6	4	4.50
4	61	16	7	4	3	3.90	116	29	13	8	5	5.37	134	34	15	9	6	5.77
5	63	16	7	4	3	3.96	129	33	15	9	6	5.67	136	34	16	9	6	5.82
6	102	26	12	7	5	5.05	126	32	14	8	6	5.61	158	40	18	10	7	6.28
7	95	24	11	6	4	4.85	193	49	22	13	8	6.94	261	66	29	17	11	8.07
8	88	22	10	6	4	4.69	139	35	16	9	6	5.89	218	55	25	14	9	7.38
9	228	57	26	15	10	7.55	349	88	39	22	14	9.34	308	77	35	20	13	8.76
10	349	88	39	22	14	9.33	519	130	58	33	21	11.38	627	157	70	40	26	12.51
11	36	9	4	3	2	2.98	546	137	61	35	22	11.68	534	134	60	34	22	11.55
Sowing date 2 (05/25/2016)																		
1	166	42	19	11	7	6.43	36	9	4	3	2	2.99	- <sup>(1)</sup>	-	-	-	-	-
2	119	30	14	8	5	5.44	105	27	12	7	5	5.12	117	30	13	8	5	5.40
3	113	29	13	8	5	5.30	165	42	19	11	7	6.40	92	23	11	6	4	4.78
4	70	18	8	5	3	4.16	128	32	15	8	6	5.64	144	36	16	9	6	5.99
5	72	18	8	5	3	4.21	173	44	20	11	7	6.57	182	46	21	12	8	6.74
6	72	18	8	5	3	4.24	112	28	13	7	5	5.28	148	37	17	10	6	6.08
7	63	16	7	4	3	3.94	205	52	23	13	9	7.15	422	106	47	27	17	10.27
8	65	17	8	5	3	4.01	363	91	41	23	15	9.52	587	147	66	37	24	12.10
9	275	69	31	18	11	8.28	572	143	64	36	23	11.96	626	157	70	40	26	12.50
10	52	13	6	4	3	3.6	680	170	76	43	28	13.03	920	230	103	58	37	15.16
Sowing date 3 (06/07/2016)																		
1	159	40	18	10	7	6.30	113	29	13	8	5	5.29	-	-	-	-	-	-
2	109	28	13	7	5	5.22	68	17	8	5	3	4.11	106	27	12	7	5	5.13
3	78	20	9	5	4	4.39	90	23	10	6	4	4.74	105	27	12	7	5	5.11
4	63	16	7	4	3	3.95	108	27	12	7	5	5.19	115	29	13	8	5	5.35
5	58	15	7	4	3	3.79	164	41	19	11	7	6.39	150	38	17	10	6	6.11
6	48	12	6	3	2	3.46	120	30	14	8	5	5.46	134	34	15	9	6	5.77
7	58	15	7	4	3	3.79	254	64	29	16	11	7.96	179	45	20	12	8	6.67
8	105	27	12	7	5	5.12	284	71	32	18	12	8.42	258	65	29	17	11	8.03
9	415	104	47	26	17	10.18	374	94	42	24	15	9.66	358	90	40	23	15	9.45
10	43	11	5	3	2	3.26	486	122	54	31	20	11.02	426	107	48	27	18	10.31

continue

continuation

Sowing date 4 (06/22/2016)																		
1	162	41	18	11	7	6.35	127	32	15	8	6	5.61	-	-	-	-	-	
2	64	16	8	4	3	3.99	42	11	5	3	2	3.22	62	16	7	4	3	3.92
3	62	16	7	4	3	3.91	76	19	9	5	4	4.35	83	21	10	6	4	4.55
4	64	16	8	4	3	3.97	122	31	14	8	5	5.52	88	22	10	6	4	4.67
5	69	18	8	5	3	4.14	81	21	9	6	4	4.49	96	24	11	6	4	4.88
6	55	14	7	4	3	3.69	161	41	18	11	7	6.34	129	33	15	9	6	5.68
7	39	10	5	3	2	3.12	152	38	17	10	7	6.16	143	36	16	9	6	5.97
8	45	12	5	3	2	3.34	243	61	27	16	10	7.78	180	45	20	12	8	6.71
9	296	74	33	19	12	8.60	308	77	35	20	13	8.76	356	89	40	23	15	9.43
10	50	13	6	4	2	3.51	506	127	57	32	21	11.24	598	150	67	38	24	12.22
Sowing date 5 (07/04/2016)																		
1	121	31	14	8	5	5.49	51	13	6	4	3	3.56	203	51	23	13	9	7.11
2	109	28	13	7	5	5.22	126	32	14	8	6	5.60	179	45	20	12	8	6.67
3	79	20	9	5	4	4.43	201	51	23	13	9	7.08	122	31	14	8	5	5.51
4	65	17	8	5	3	4.02	136	34	16	9	6	5.81	110	28	13	7	5	5.23
5	60	15	7	4	3	3.84	178	45	20	12	8	6.67	201	51	23	13	9	7.08
6	49	13	6	4	2	3.49	229	58	26	15	10	7.56	242	61	27	16	10	7.77
7	55	14	7	4	3	3.70	264	66	30	17	11	8.12	324	81	36	21	13	8.99
8	127	32	15	8	6	5.63	364	91	41	23	15	9.53	266	67	30	17	11	8.14
9	533	134	60	34	22	11.54	501	126	56	32	21	11.19	579	145	65	37	24	12.03
10	46	12	6	3	2	3.39	886	222	99	56	36	14.88	580	145	65	37	24	12.04

(1) It was not possible to calculate the sample size because there was no variance in the data (plants presented only one stem).

With the option of sampling 100 plants, the estimation error for the BRS Progresso cultivar would be at most  $\pm 7.41$ ,  $11.63$ , and  $13.61\%$  of the mean, respectively, for traits PH, NL, and NS (Table 5). Meanwhile for the Temprano cultivar, sampling the same number of plants, there would be maximum estimation errors of  $\pm 11.54$ ,  $14.88$ , and  $15.16\%$  of the mean, respectively, for the traits PH, NL, and NS (Table 6).

There was a requirement of smaller sample sizes for PH trait in evaluations at the end-of-cycle (flowering stage) and greater sample sizes for NL and NS traits (Tables 5, 6). The smaller number of plants for sampling plant height also was found in end-of-cycle evaluations in *Crotalaria spectabilis* (TOEBE et al., 2017). In general, sample sizes were smaller at sowing dates 3 and 4, indicating that the plants from these sowing dates are more homogeneous. In black oat, sample sizes for intermediate evaluation times were greater than in the initial and final evaluation times (CARGNELUTTI FILHO et

al., 2015). With the variability found between the traits evaluated in the rye crop, it is evident that the volume of data collected was representative for the crop. Consequently, the sample size to estimate the mean of these traits can be taken as reference for future studies with this crop.

## Conclusion

There is variability of sample size estimate between traits plant height, number of leaves, and number of stems, between cultivars BRS Progresso and Temprano, between sowing dates, and between evaluation times. Larger sample sizes were required for the Temprano cultivar than the BRS Progresso cultivar. For the BRS Progresso cultivar, 25, 61, and 83 plants were required to estimate mean plant height, number of leaves, and number of stems, respectively, with a maximum estimation error of  $15\%$ . For the Temprano cultivar, 60, 99, and 103 plants were required to estimate mean plant height,

number of leaves, and number of stems, respectively, with a maximum estimation error of 15%.

## Acknowledgments

We thank the Brazilian National Council for Scientific and Technological Development (CNPq) and the Coordination for the Improvement of Higher Education Personnel (CAPES) for granting scholarships. Our acknowledgement to the Embrapa Trigo, for donating the rye, cultivar BRS Progreso seeds that were used in this study, as well as scholarship students for their help in data collection.

## References

- BAIER, A. C. *Centeio*. Passo Fundo: EMBRAPA, 1994. 29 p. (Documento, 15).
- BUSSAB, W. O.; MORETTIN, P. A. *Estatística básica*. São Paulo: Saraiva, 2013. 548 p.
- CAMPOS, H. *Estatística experimental não-paramétrica*. 4. ed. Piracicaba: Escola Superior de Agricultura Luiz de Queiroz da Universidade de São Paulo, 1983. 349 p.
- CARGNELUTTI FILHO, A.; EVANGELISTA, D. H. R.; GONÇALVES, E. C. P.; STORCK, L. Tamanho de amostra de caracteres de genótipos de soja. *Ciência Rural*, Santa Maria, v. 39, n. 4, p. 983-991, 2009.
- CARGNELUTTI FILHO, A.; LOPES, S. J.; BRUM, B.; SILVEIRA, T. R.; TOEBE, M.; STORCK, L. Tamanho de amostra de caracteres em híbridos de mamoneira. *Ciência Rural*, Santa Maria, v. 40, n. 2, p. 280-287, 2010.
- CARGNELUTTI FILHO, A.; TOEBE, M.; ALVES, B. M.; BURIN, C.; SANTOS, G. O.; FACCO, G.; NEU, I. M. M. Dimensionamento amostral para avaliar caracteres morfológicos e produtivos de aveia preta em épocas de avaliação. *Ciência Rural*, Santa Maria, v. 45, n. 1, p. 9-13, 2015.
- COMPANHIA NACIONAL DE ABASTECIMENTO - CONAB. Acompanhamento da safra brasileira de grãos, v. 4 - Safra 2016/17, n. 8 - Oitavo levantamento. Brasília: Conab, 2017. Disponível em: <[http://www.conab.gov.br/OlalaCMS/uploads/arquivos/17\\_05\\_12\\_10\\_37\\_57\\_boletim\\_graos\\_maior\\_2017.pdf](http://www.conab.gov.br/OlalaCMS/uploads/arquivos/17_05_12_10_37_57_boletim_graos_maior_2017.pdf)>. Disponível em: 15 fev. 2018.
- CRUZ, C. D. GENES - a software package for analysis in experimental statistics and quantitative genetics. *Acta Scientiarum. Agronomy*, Maringá, v. 35, n. 3, p. 271-276, 2013.
- FACCO, G.; CARGNELUTTI FILHO, A.; ALVES, B. M.; BURIN, C.; SANTOS, G. O.; KLEINPAUL, J. A.; NEU, I. M. M. Sample size for estimating average productive traits of pigeon pea. *Ciência Rural*, Santa Maria, v. 46, n. 4, p. 619-625, 2016.
- FACCO, G.; CARGNELUTTI FILHO, A.; LÚCIO, A. D.; SANTOS, G. O.; STEFANELLO, R. B.; ALVES, B. M.; BURIN, C.; NEU, I. M. M.; KLEINPAUL, J. A. Sample size for morphological traits of pigeon pea. *Semina: Ciências Agrárias*, Londrina, v. 36, n. 6, p. 4151-4164, 2015.
- HELDWEIN, A. B.; BURIOL, G. A.; STRECK, N. A. O clima de Santa Maria. *Ciência & Ambiente*, Santa Maria, v. 38, n. 1, p. 43-58, 2009.
- LIN, M.; LUCAS, H. C.; SHMUELI, G. Too big to fail: large samples and the p-value problem. *Information Systems Research*, Catonsville, v. 24, n. 4, p. 906-917, 2013.
- PANTOJA, J. L.; WOLI, K. P.; SAWYER, J. E.; BARKER, D. W. Winter rye cover crop biomass production, degradation, and nitrogen recycling. *Agronomy Journal*, Madison, v. 108, n. 2, p. 841-853, 2016.
- SANTOS, H. G.; JACOMINE, P. K. T.; ANJOS, L. H. C.; OLIVEIRA, V. A.; LUMBRERAS, J. F.; COELHO, M. R.; ALMEIDA, J. A.; CUNHA, T. J. F.; OLIVEIRA, J. B. *Sistema brasileiro de classificação de solos*. 3. ed. Brasília: Embrapa, 2013. 353 p.
- STORCK, L.; LOPES, S. J.; CARGNELUTTI FILHO, A.; MARTINI, L. F. D.; CARVALHO, M. P. Sample size for single, double and three-way hybrid corn ear traits. *Scientia Agrícola*, Piracicaba, v. 64, n. 1, p. 30-35, 2007.
- TEODORO, P. E.; SILVA JUNIOR, C. A.; RIBEIRO, L. P.; SILVA, F. A.; CORRÊA, C. C. G.; ZANUNCIO, A. S.; TORRES, F. E. Sample dimension for estimation of biomass and yield of sunn (*Crotalaria juncea* L.) and showy rattlebox (*C. spectabilis* Roth.). *Journal of Agronomy*, Faisalabad, v. 14, n. 2, p. 98-101, 2015.
- TOEBE, M.; BANDEIRA, C. T.; FORTES, S. K. G.; CARVALHO, J. O.; TARTAGLIA, F. L.; TAMBARA, A. L.; MELO, P. J. Dimensionamento amostral e associação linear entre caracteres de *Crotalaria spectabilis*. *Bragantia*, Campinas, v. 76, n. 1, p. 45-53, 2017.
- ZIECH, A. R. D.; CONCEIÇÃO, P. C.; LUCHESE, A. V.; BALIN, N. M.; CANDIOTTO, G.; GARMUS, T. G. Proteção do solo por plantas de cobertura de ciclo hibernar na região Sul do Brasil. *Pesquisa Agropecuária Brasileira*, Brasília, v. 50, n. 5, p. 374-382, 2015.